

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁵ : C07D 401/12, A61K 31/42, 31/445, C07D 409/12, 413/12, 417/12, 487/08, 261/20, 263/56, 275/04, 277/64, 277/66, 333/54</p>	<p>A1</p>	<p>(11) International Publication Number: WO 94/13663 (43) International Publication Date: 23 June 1994 (23.06.94)</p>
<p>(21) International Application Number: PCT/US93/09407 (22) International Filing Date: 7 October 1993 (07.10.93) (30) Priority Data: 07/988,653 10 December 1992 (10.12.92) US (60) Parent Application or Grant (63) Related by Continuation US 07/988,653 (CIP) Filed on 10 December 1992 (10.12.92) (71) Applicant (for all designated States except US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HOWARD, Harry, R. [US/US]; 272 Westwoods Terrace, Bristol, CT 06010 (US). O'NEILL, Brian, T. [US/US]; 1526 Essex Road, Westbrook, CT 06498 (US). (74) Agents: RICHARDSON, Peter et al.; Pfizer Inc., 4G235 East 42nd Street, New York, NY 10017 (US).</p>		<p>(81) Designated States: AU, CA, JP, KR, NO, NZ, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report.</p>
<p>(54) Title: AMINOMETHYLENE SUBSTITUTED NON-AROMATIC HETEROCYCLES AND USE AS SUBSTANCE P ANTAGONISTS</p> <div style="display: flex; justify-content: space-around; align-items: flex-start;"> <div data-bbox="243 1155 730 1323"> <p style="text-align: center;">(Ia)</p> </div> <div data-bbox="925 1155 1396 1365"> <p style="text-align: center;">(Ib)</p> </div> </div> <p>(57) Abstract</p> <p>The present invention relates to novel aminomethylene substituted non-aromatic heterocycles and, specifically, to compounds of formula (Ia) or (Ib) wherein W, R¹, R², R³, A, X', Y' and Z' are as defined in the specification, and to intermediates used in the synthesis of such compounds. The novel compounds of formulae (Ia) and (Ib) are useful in the treatment of inflammatory and central nervous system disorders, as well as other disorders.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

AMINOMETHYLENE SUBSTITUTED NON-AROMATIC HETEROCYCLES AND USE
AS SUBSTANCE P ANTAGONISTS

Background of the Invention

The present invention relates to novel aminomethylene substituted non-aromatic heterocycles, pharmaceutical compositions comprising such compounds and the use of such compounds in the treatment and prevention of inflammatory and central nervous system disorders, as well as several other disorders. The pharmaceutically active compounds of this invention are substance P receptor antagonists. This invention also relates to novel intermediates used in the synthesis of such substance P receptor antagonists.

Substance P is a naturally occurring undecapeptide belonging to the tachykinin family of peptides, the latter being named because of their prompt stimulatory action on smooth muscle tissue. More specifically, substance P is a pharmacologically active neuropeptide that is produced in mammals and possesses a characteristic amino acid sequence that is illustrated by D. F. Veber et al. in U.S. Patent No. 4,680,283.


The following references refer, collectively, to quinuclidine, piperidine, and azanorbornane derivatives and related compounds that exhibit activity as substance P receptor antagonists: United States Patent 5,162,339, which issued on November 11, 1992; United States Patent Application 724,268, filed July 1, 1991; PCT Patent Application PCT/US 91/02853, filed April 25, 1991; PCT Patent Application PCT/US 91/03369, filed May 14, 1991; PCT Patent Application PCT/US 91/05776, filed August 20, 1991; PCT Patent Application PCT/US 92/00113, filed January 17, 1992; PCT Patent Application PCT/US 92/03571, filed May 5, 1992; PCT Patent Application PCT/US 92/03317, filed April 28, 1992; PCT Patent Application PCT/US 92/04697, filed June 11, 1992; United States Patent Application 766,488, filed September 26, 1991; United States Patent Application 790,934, filed November 12, 1991; PCT Patent Application PCT/US 92/04002, filed May 19, 1992; Japanese Patent

-2-

Application 065337/92, filed March 23, 1992; and United States Patent Application 932,392, filed August 19, 1992.


Summary of the Invention

The present invention relates to compounds of the
5 formula



10

or



15

wherein A is a ring system selected from phenyl, naphthyl,
20 thienyl, quinoliny and indoliny, and wherein the side chain containing NR^2R^3 is attached to a carbon atom of ring system A;

W is hydrogen, (C_1-C_6) alkyl optionally substituted with from one to three fluorine atoms, $-S(O)_v-(C_1-C_6)$ alkyl wherein
25 v is zero, one or two, halo, benzyloxy or (C_1-C_6) alkoxy optionally substituted with from one to three fluorine atoms;

R^1 is a 4, 5 or 6 membered heterocyclic ring containing from one to three heteroatoms selected from oxygen, nitrogen
30 and sulfur (e.g., thiazolyl, azetidyl, pyrrolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, isothiazolyl, imidazolyl, isoxazolyl, oxazolyl, pyridyl, pyrimidinyl, pyrazolyl or thiophenyl), wherein said heterocyclic ring may contain from zero to three double bonds and may optionally
35 be substituted with one or more substituents, preferably one or two substituents, independently selected from (C_1-C_6) alkyl optionally substituted with from one to three fluorine

-3-

atoms and (C₁-C₆) alkoxy optionally substituted with from one to three fluorine atoms;

the dotted lines in formula Ib indicate that one of the X'-Y' and Y'-Z' bonds may optionally be a double bond;

5 X' is selected from =CH-, -CH₂-, -O-, -S-, -SO-, -SO₂-, -N(R⁴)-, -NH-, =N-, -CH[(C₁-C₆)alkyl]-, =C[(C₁-C₆)alkyl]-, -CH(C₆H₅)- and =C(C₆H₅)-;

Y' is selected from C=O, C=NR⁴, C=S, =CH-, -CH₂-, =C[(C₁-C₆)alkyl]-, -CH[(C₁-C₆)alkyl]-, =C(C₆H₅)-, -CH(C₆H₅)-, =N-,
10 -NH-, -N(R⁴)-, =C(halo)-, =C(OR⁴)-, =C(SR⁴)-, =C(NR⁴)-, -O-, =C(CF₃)-, =C(CH₂C₆H₅)-, -S- and SO₂, wherein the phenyl moieties of said =C(C₆H₅)- and -CH(C₆H₅)- may optionally be substituted with from one to three substituents independently selected from trifluoromethyl and halo, and
15 wherein the alkyl moieties of said =[(C₁-C₆)alkyl]- and -CH[(C₁-C₆)alkyl]- may optionally be substituted with from one to three fluorine atoms;

Z' is selected from =CH-, -CH₂-, =N-, -NH-, -S-, -N(R⁴)-, =C(C₆H₅)-, -CH(C₆H₅)-, =C[(C₁-C₆)alkyl]- and -CH[(C₁-
20 C₆)alkyl]-;

or X', Y' and Z', together with the two carbon atoms shared between the benzo ring and the X'Y'Z' ring, form a fused pyridine or pyrimidine ring;

R² is hydrogen or -CO₂(C₁-C₁₀)alkyl;

25 R³ is selected from

-5-

wherein R^6 and R^{10} are independently selected from furyl, thienyl, pyridyl, indolyl, biphenyl and phenyl, wherein said phenyl may optionally be substituted with one or two substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms, (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms, carboxy, benzyloxycarbonyl and (C_1-C_3) alkoxy-carbonyl;


R^4 is (C_1-C_6) alkyl or phenyl;

10 R^7 is selected from (C_3-C_4) branched alkyl, (C_5-C_6) branched alkenyl, (C_5-C_7) cycloalkyl, and the radicals named in the definition of R^6 ;

R^8 is hydrogen or (C_1-C_6) alkyl;

15 R^9 and R^{19} are independently selected from phenyl, biphenyl, naphthyl, pyridyl, benzhydryl, thienyl and furyl, and R^9 and R^{19} may optionally be substituted with from one to three substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms and (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms;

20 Y is $(CH_2)_l$, wherein l is an integer from one to three, or Y is a group of the formula



Z is oxygen, sulfur, amino, (C_1-C_3) alkylamino or $(CH_2)_n$,
 30 wherein n is zero, one or two;
 x is zero, one or two;
 y is zero, one or two;
 z is three, four or five;
 o is two or three;
 35 p is zero or one;
 r is one, two or three;

-6-

the ring containing $(CH_2)_z$ may contain from zero to three double bonds, and one of the carbon atoms of $(CH_2)_z$ may optionally be replaced by oxygen, sulfur or nitrogen;

R^{11} is thienyl, biphenyl or phenyl optionally substituted with one or two substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms and (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms;



10 X is $(CH_2)_q$ wherein q is an integer from 1 to 6, and wherein any one of the carbon-carbon single bonds in said $(CH_2)_q$ may optionally be replaced by a carbon-carbon double bond, and wherein any one of the carbon atoms of said $(CH_2)_q$ may optionally be substituted with R^{14} , and wherein any one
15 of the carbon atoms of said $(CH_2)_q$ may optionally be substituted with R^{15} ;

m is an integer from 0 to 8, and any one of the carbon-carbon single bonds of $(CH_2)_m$, wherein both carbon atoms of such bond are bonded to each other and to another
20 carbon atom of the $(CH_2)_m$ chain, may optionally be replaced by a carbon-carbon double bond or a carbon-carbon triple bond, and any one of the carbon atoms of said $(CH_2)_m$ may optionally be substituted with R^{17} ;

R^{12} is a radical selected from hydrogen, (C_1-C_6) straight or branched alkyl, (C_3-C_7) cycloalkyl wherein one of the carbon atoms may optionally be replaced by nitrogen, oxygen or sulfur; aryl selected from biphenyl, phenyl, indanyl and naphthyl; heteroaryl selected from thienyl, furyl, pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl and quinolyl; phenyl- (C_2-C_6) alkyl, benzhydryl and benzyl, wherein the point of attachment on R^{12} is a carbon atom unless R^{12} is hydrogen, and wherein each of said aryl and heteroaryl groups and the phenyl moieties of said benzyl, phenyl- (C_2-C_6) alkyl and benzhydryl may optionally be
30 substituted with one or more substituents independently selected from halo, nitro, (C_1-C_{10}) alkyl optionally
35

-7-


substituted with from one to three fluorine atoms, (C₁-C₁₀)
alkoxy optionally substituted with from one to three
fluorine atoms, amino, hydroxy-(C₁-C₆)alkyl,
(C₁-C₆)alkoxy-(C₁-C₆)alkyl, (C₁-C₆)-alkylamino,




R¹³ is hydrogen, phenyl or (C₁-C₆)alkyl;

or R¹² and R¹³, together with the carbon to which they
are attached, form a saturated carbocyclic ring having from
30 3 to 7 carbon atoms wherein one of said carbon atoms that is
neither the point of attachment of the spiro ring nor
adjacent to such point of attachment may optionally be
replaced by oxygen, nitrogen or sulfur;

R¹⁴ and R¹⁵ are each independently selected from
35 hydrogen, hydroxy, halo, amino, oxo (=O), cyano, hydroxy-(C₁-
C₆)alkyl, (C₁-C₆)alkoxy-(C₁-C₆)alkyl, (C₁-C₆)alkylamino,



-8-



5 $(C_1-C_6)\text{alkyl}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-$, $(C_1-C_6)\text{alkyl}-\overset{\overset{\text{O}}{\parallel}}{\text{C}}-(C_1-C_6)\text{alkyl}-$, and the radicals
 10 set forth in the definition of R^{12} ;

15 R^{16} is $\text{NH}\overset{\overset{\text{O}}{\parallel}}{\text{C}}\text{R}^{18}$, $\text{NHCH}_2\text{R}^{18}$, SO_2R^{18} , CO_2H or one of the
 radicals set forth in any of the definitions of R^{12} , R^{14} and
 R^{15} ;

R^{17} is oximino ($=\text{NOH}$) or one of the radicals set forth
 in any of the definitions of R^{12} , R^{14} and R^{15} ; and

R^{18} is $(C_1-C_6)\text{alkyl}$, hydrogen, phenyl or phenyl $(C_1-C_6)\text{alkyl}$;

20 with the proviso that (a) when m is 0, one of R^{16} and R^{17}
 is absent and the other is hydrogen, (b) when R^3 is a group
 of the formula VIII, R^{14} and R^{15} cannot be attached to the
 same carbon atom, (c) when R^{14} and R^{15} are attached to the
 same carbon atom, then either each of R^{14} and R^{15} is
 25 independently selected from hydrogen, fluoro, $(C_1-C_6)\text{alkyl}$,
 hydroxy- $(C_1-C_6)\text{alkyl}$ and $(C_1-C_6)\text{alkoxy-}(C_1-C_6)\text{alkyl}$, or R^{14} and
 R^{15} , together with the carbon to which they are attached,
 form a (C_3-C_6) saturated carbocyclic ring that forms a spiro
 compound with the nitrogen-containing ring to which they are
 30 attached; (d) R^{12} and R^{13} can not both be hydrogen, and (e)
 when R^{14} or R^{15} is attached to a carbon atom of X or $(\text{CH}_2)_n$,
 that is adjacent to the ring nitrogen, then R^{14} or R^{15} ,
 respectively, must be a substituent wherein the point of
 attachment is a carbon atom.

35 The present invention also relates to the
 pharmaceutically acceptable acid addition and base salts of
 compounds of the formulae Ia and Ib (hereinafter referred
 to, collectively, as compounds of the formula I). The acids
 which are used to prepare the pharmaceutically acceptable
 40 acid addition salts of the aforementioned base compounds of

-9-

this invention are those which form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, acetate, lactate, citrate, acid citrate, tartrate, bitartrate, succinate, maleate, fumarate, gluconate, saccharate, benzoate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and pamoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)]salts. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula I. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium calcium and magnesium, etc.

The fused bicyclic nucleus of compounds of the formula Ib to which W and the $-CH_2NR^2R^3$ sidechain are attached may be, but is not limited to one of the following groups: benzoxazolyl, benzthiazolyl, benzimidazolyl, benzisoxazolyl, benzoisothiazolyl, indazolyl, indolyl, isoquinolinyl, benzofuryl, benzothienyl, oxindolyl, benzoxazolinonyl, benzthiazolinonyl, benzimidazolinonyl, benzimidazoliniminy, dihydrobenzothienyl-S,S-dioxide, benztriazolyl, benzthiadiazolyl, benzoxadiazolyl, and quinazolinyl.

The term "halo", as used herein, unless otherwise indicated, includes chloro, fluoro, bromo and iodo.

The term "alkyl", as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.

The term "alkoxy", as used herein, includes O-alkyl groups wherein "alkyl" is defined as above.

The term "one or more substituents," as used herein, includes from one to the maximum number of substituents possible based on the number of available bonding sites.

-10-

Preferred compounds of this invention include those compounds of the formula I wherein the substituents at positions "2" and "3" of the nitrogen containing ring of R³ are in a cis configuration. When R³ is a group of the formula VII or VIII, "a cis configuration", as used herein, means that the non-hydrogen substituent at position "3" is cis to R¹².

Other preferred compounds of this invention include those compounds of the formula Ia wherein R³ is a group of the formula III, VII or IX; R² is hydrogen; A is phenyl or indolyl; W is (C₁-C₃)alkoxy optionally substituted with from one to five fluorine atoms; and R¹ is thiazolyl, imidazolyl, thiadiazolyl, pyrrolyl, oxazolyl, pyridyl, pyrimidinyl, pyrazolyl or thiophenyl, and R¹ may optionally be substituted with one or two (C₁-C₃) alkyl moieties.

Other preferred compounds of this invention include those compounds of the formula Ib wherein R³ is a group of the formula III, VII or IX; R² is hydrogen; the fused bicyclic ring system to which W and the -CH₂NR²R³ sidechain are attached is benzoxazolyl, benzisoxazolyl, benzthiazolyl, benzthiophenyl or benzimidazolyl; and W is (C₁-C₆)alkoxy optionally substituted with from one to five fluorine atoms.

More preferred compounds of this invention are the foregoing preferred compounds wherein: (a) R³ is a group of the formula III and R⁹ is benzhydryl; (b) R³ is a group of the formula VII, R¹² is phenyl, each of R¹³, R¹⁴, R¹⁵ and R¹⁶ is hydrogen, m is zero and X is -(CH₂)₃-; or (c) R³ is a group of the formula IX, r is two and R¹⁹ is benzhydryl.

Other more preferred compounds of this invention are those compounds of the formula Ia wherein: (a) R³ is a group of the formula III wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, R⁹ is benzhydryl and A is phenyl; or (b) R³ is a group of the formula VII wherein R¹² and the substituent at position "3" of the nitrogen containing ring are in the cis configuration, A is phenyl, R¹² is phenyl, each of R², R¹³, R¹⁴,

-11-

R¹⁵ and R¹⁶ is hydrogen, m is zero, W is methoxy or isopropoxy, X is -(CH₂)₃- and R¹ is thiazolyl, imidazolyl, pyrrolyl, oxazolyl, pyridyl, pyrimidinyl, pyrazolyl, thiophenyl or thiadiazolyl.

5 Other more preferred compounds of this invention are those compounds of the formula Ib wherein R³ is a group of the formula IX wherein the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, R¹⁹ is benzhydryl, r is two and the fused
10 bicyclic ring system to which W and the -CH₂NR²R³ sidechain are attached is benzisoxazolyl or benzthiazolyl.

Especially preferred compounds of the formula Ib are those wherein R³ is a group of the formula IX, R¹⁹ is benzhydryl, the fused bicyclic ring system to which W and
15 the -CH₂NR²R³ sidechain are attached is benzisoxazolyl, and W is methoxy.

Other especially preferred compounds of the formula Ib are those wherein R³ is a group of the formula VII, R¹² is phenyl, each of R¹³, R¹⁴, R¹⁵ and R¹⁶ is hydrogen, m is zero, X
20 is -(CH₂)₃-, and the fused bicyclic ring system to which W and the -CH₂NR²R³ sidechain are attached is benzothiazolyl, benzoxazolyl, benzthiophenyl or benzimidazolyl.

Especially preferred compounds of the formula Ia are those wherein R³ is a group of the formula VII, each of R¹³,
25 R¹⁴, R¹⁵ and R¹⁶ is hydrogen, m is zero, X is -(CH₂)₃-, A is phenyl, W is methoxy, and R¹ is selected from thiazolyl, imidazolyl, thiadiazolyl, pyridyl, pyrimidinyl, pyrazolyl, thiophenyl and isoxazolyl.

Specific preferred compounds of the formula I include
30 the following:

(2S,3S)-3-[2-methoxy-5-(2-thiazolyl)benzyl]amino-2-phenylpiperidine;

(2S,3S)-3-[5-(2-imidazolyl)-2-methoxybenzyl]amino-2-phenylpiperidine;

35 (2S,3S)-3-[2-methoxy-5-(2-oxopyrrolidinyl)benzyl]amino-2-phenylpiperidine;

-12-

- (2S,3S)-3-[2-methoxy-5-(4-methyl-2-thiazolyl)benzyl]-amino-2-phenylpiperidine;
- (2S,3S)-3-[2-methoxy-5-(1,2,3-thiadiazol-4-yl)benzyl]-amino-2-phenylpiperidine;
- 5 (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-[5-(2,5-dimethyl-pyrrol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-3-[2-methoxy-5-(5-oxazolyl)benzyl]amino-2-phenylpiperidine;
- 10 (2S,3S)-(6-methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;
- (2S,3S)-(6-methoxy-2-cyclopropyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- 15 (2S,3S)-(6-methoxy-2-tert-butyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-isopropoxyoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-isopropoxyoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- 20 (2S,3S)-(6-trifluoromethoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-methoxy-2-methyl-benzoxazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- 25 (1SR,2SR,3SR,4RS)-3-[6-methoxy-3-methylbenzisoxazol-5-yl]methylamino-2-benzhydrylazanorbornane;
- (2S,3S)-(2-methoxy-5-pyridin-2-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(2-methoxy-5-pyrimidin-2-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;
- 30 (2S,3S)-(2-methoxy-5-pyridin-3-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-[2-methoxy-5-(6-methylpyridin-2-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;
- 35 (2S,3S)-[5-(3,5-dimethylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;

-13-

(2S,3S)-[2-methoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[2-isopropoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;

5 (2S,3S)-[5-(3,5-diisopropylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[5-(3,5-dimethylthiophen-2-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine; and

10 (2S,3S)-(6-methoxy-2,3-dimethyl-benzo[b]thiophen-7-ylmethyl)-(2-phenylpiperidin-3-yl)amine.

Other compounds of the formula I include the following:

(2S,3S)-(6-methoxy-3-methyl-benzo[d]isoxazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;

15 (1SR,2SR,3SR,4RS)-(2-benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-amine;

(2S,3S)-(6-methoxy-benzoxazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;

20 (2S,3S)-(6-methoxy-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;

(2S,3S)-5-methoxy-1-methyl-6-(2-phenylpiperidin-3-ylaminomethyl)-1,3-dihydro-indol-2-one;

(2S,3S)-6-methoxy-3-methyl-5-(2-phenylpiperidin-3-ylaminomethyl)-3H-benzoxazol-2-one;

25 (2S,3S)-6-methoxy-3-methyl-5-(2-phenylpiperidin-3-ylaminomethyl)-3H-benzothiazol-2-one;

(2S,3S)-5-methoxy-1,3-dimethyl-6-(2-phenylpiperidin-3-ylaminomethyl)-1,3-dihydro-benzimidazol-2-one;

30 (2S,3S)-(6-methoxy-3-methyl-3H-benzotriazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-(2-methoxy-5-[1,2,3]thiadiazol-4-yl-benzyl)-(2-phenyl-1-azabicyclo[2.2.2]oct-3-yl)amine;

(2S,3S)-(2-methoxy-5-[1,2,3]thiadiazol-4-yl-benzyl)-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-yl)amine;

35 (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenyl-1-azabicyclo[2.2.2]oct-3-yl)amine;

-14-

- (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-yl)amine;
- (2S,3S)-(2-methoxy-5-thiazol-2-yl-benzyl)-(2-benzhydryl-1-azabicyclo[2.2.2]oct-3-yl)amine;
- 5 (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenyl-1-azabicyclo[2.2.1]hept-3-yl)amine;
- (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)amine;
- (2S,3S)-(2-methoxy-5-[1,2,4]triazol-4-yl-benzyl)-(2-phenylpiperidin-3-yl)amine;
- 10 (2S,3S)-(2-methoxy-5-[1,2,4]triazol-1-yl-benzyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(2-methoxy-5-thiazol-2-ylbenzyl)-(2-phenyl-decahydroquinolin-3-yl)amine;
- 15 (2S,3S)-(2-methoxy-5-thiazol-2-ylbenzyl)-(2-phenyl-octahydro-indol-3-yl)amine;
- (2S,3S)-(2-methoxy-5-oxazol-4-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-methoxy-2-(2-propyl)-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;
- 20 (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-phenyl-benzothiazol-5-ylmethyl)amine;
- (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-cyclopropyl-benzothiazol-5-ylmethyl)amine;
- 25 (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-tert-butyl-benzothiazol-5-ylmethyl)amine;
- (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-(2-propyl)-benzothiazol-5-ylmethyl)amine;
- 30 (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-(2-propyl)-benzothiazol-5-ylmethyl)amine;
- (1SR, 2SR, 3SR, 4RS)-(2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl)-(6-isopropoxyoxy-2-phenyl-benzothiazol-5-ylmethyl)amine;
- 35

-15-

(1SR, 2SR, 3SR, 4RS) - (2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl) - (6-isopropoxyoxy-2-methylbenzothiazol-5-ylmethyl)amine;


(1SR, 2SR, 3SR, 4RS) - (2-benzhydryl-1-azabicyclo[2.2.1]hept-3-yl) - (6-trifluoromethoxy-2-methylbenzothiazol-5-ylmethyl)amine;

(6-methoxy-1-oxa-2,3-diazainden-5-ylmethyl) - (2-phenylpiperidin-3-yl)amine; and


(6-methoxy-2-methyl-1H-benzoimidazol-5-ylmethyl) - (2-phenylpiperidine-3-yl)amine.

The present invention also relates to compounds of the formulae

-16-




XI




XII

10




XI-A

and



XII-A

wherein ring A, R¹, R³, W, X', Y' and Z' are defined as
20 above. It also relates to compounds of the formula



XVIII

25

wherein X' is -S- or -O-, and each of Y' and Z' is,
independently, =N-, =CH-, =C[(C₁-C₆)alkyl]- or =C(C₆H₅)-,
wherein the alkyl moiety of said =C[(C₁-C₆)alkyl]- may
optionally be substituted with from one to three fluorine
atoms and the phenyl moiety of said =C(C₆H₅)- may optionally
30 be substituted with from one to three substituents
independently selected from halo and trifluoromethyl, with
the proviso that Y' and Z' can not both be =N-, and R²² is
methyl, ethyl, n-propyl, isopropyl, t-butyl,
35 trifluoromethyl, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or benzyl.
Compounds of the formulae XI, XII, XI-A, XII-A and XVIII are

-17-

intermediates in the synthesis of compounds of the formulae Ia and Ib.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, urinary incontinence, gastrointestinal disorders such as emesis and colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such condition, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, urinary incontinence, gastrointestinal disorders such as emesis and colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as

-18-

shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic
5 neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising administering to said mammal an amount of a compound of the formula I, or a
10 pharmaceutically acceptable salt thereof, effective in treating or preventing such condition.

The present invention also relates to a pharmaceutical composition for antagonizing the effects of substance P in a mammal, including a human, comprising a substance P
15 antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of antagonizing the effects of substance P in a mammal,
20 including a human, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a
25 mammal, including a human, resulting from an excess of substance P, comprising a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

30 The present invention also relates to a method of treating or preventing a disorder in a mammal, including a human, resulting from an excess of substance P, comprising administering to said mammal a substance P antagonizing amount of a compound of the formula I, or a pharmaceutically
35 acceptable salt thereof.

The present invention also relates to a pharmaceutical composition for treating or preventing a condition selected

-19-

from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, urinary incontinence, gastrointestinal disorders such as emesis and colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in a mammal, including a human, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a condition selected from the group consisting of inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, urinary incontinence, gastrointestinal disorders such as emesis and colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as

-20-

Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis in
5 a mammal, including a human, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical
10 composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable
15 salt thereof, effective in antagonizing the effect of substance P at its receptor site, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of treating or preventing a disorder in mammal, including a
20 human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in
25 antagonizing the effect of substance P at its receptor site.

The present invention also relates to a pharmaceutical composition for treating or preventing a disorder in a mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance
30 P mediated neurotransmission, comprising an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder, and a pharmaceutically acceptable carrier.

The present invention also relates to a method of
35 treating or preventing a disorder in mammal, including a human, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated

-21-

neurotransmission, comprising administering to said mammal an amount of a compound of the formula I, or a pharmaceutically acceptable salt thereof, effective in treating or preventing such disorder.


5 The compounds of the formula I have chiral centers and therefore exist in different enantiomeric forms. This invention relates to all optical isomers and all stereoisomers of compounds of the formula I, and mixtures thereof.

10 Detailed Description of the Invention



The compounds of the formula I may be prepared as described in the following reaction schemes and discussion. Unless otherwise indicated, ring A, W, R¹, R², R³, R⁴, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², R¹³, R¹⁴, R¹⁵, R¹⁶, R¹⁷, R¹⁸, R¹⁹, R²², X, Z, Y,
15 X', Y', Z', m, n, o, p, q, r, x, y, and z, and structural formulas Ia, Ib, II, III, IV, V, VI, VII, VIII, IX, XI, XII, XI-A and XII-A in the reaction schemes and discussion that follow are defined as above.

As indicated above, compounds of the formulae Ia and Ib
20 are referred to, collectively, as "compounds of the formula I".

-22-

Scheme 1

-23-

Scheme 2Scheme 3

-24-

Scheme 4

5

10


15

20

25

30


35




-25-

Scheme 4
(continued)

5



+




10

XXIV


XXV



15



+




20

XXVI

XXVII



25



30

XXVIII


35

-26-

Scheme 1 illustrates the preparation of compounds of the formula Ia from starting materials of the formula X wherein G is hydrogen, hydroxy, chloro, bromo or (C₁-C₆)alkoxy.

5 Referring to scheme 1, a compound of the formula X wherein G is hydrogen may be converted directly into the corresponding compound of the formula I by reacting it with a compound of the formula NH₂R³ in the presence of a reducing agent. Reducing agents that may be used include sodium
10 cyanoborohydride, sodium triacetoxyborohydride, sodium borohydride, hydrogen and a metal catalyst, zinc and hydrochloric acid, and formic acid. This reaction is typically conducted in a reaction inert solvent at a temperature from about 0°C to about 150°C. Suitable
15 reaction inert solvents include lower alcohols (e.g., methanol, ethanol and isopropanol), 1,2-dichloroethane, acetic acid and tetrahydrofuran (THF). Preferably, the solvent is acetic acid, the temperature is about 25°C, the reducing agent is sodium triacetoxyborohydride, and the
20 reaction is conducted in the presence of a dehydrating agent such as molecular sieves.

Alternatively, the reaction of a compound of the formula X with a compound of the formula NH₂R³ may be carried out in the presence of a dehydrating agent or by using an
25 apparatus designed to remove azeotropically the water generated, to produce an imine of the formula



XI

35 which is then reacted with a reducing agent as described above, preferably with sodium triacetoxyborohydride in an acetic acid or 1,2-dichloroethane solvent at about room

-27-

temperature. The preparation of the imine XI is generally carried out in a reaction inert solvent such as benzene, xylene or toluene, preferably toluene, at a temperature from about 25°C to about 110°C, preferably at about the reflux temperature of the solvent. Suitable dehydrating agents/solvent systems include titanium tetrachloride/dichloromethane, titanium isopropoxide/dichloromethane and molecular sieves. Titanium tetrachloride/dichloromethane is preferred.

10 Compounds of the formula X wherein G is hydroxy, chloro, bromo or (C₁-C₆)alkoxy may be converted into the corresponding compounds of formula XII having the desired R³ group by reacting them with the appropriate compound of the formula NH₂R³ under conditions that will be obvious to those skilled in the art, and then reducing the resulting amides to yield the desired compounds having formula I wherein R² is hydrogen. When G is hydroxy, the compound of formula X is reacted with NH₂R³ in the presence of an activating agent. Appropriate activating agents include carbonyldiimidazole, chloroformates such as isobutyl chloroformate, diethylphosphoryl cyanide and dicyclohexylcarbodiimide. Carbonyldiimidazole is preferred. This reaction is generally conducted at a temperature from about 0°C to about 50°C, preferably at about 25°C, in an inert solvent such as chloroform, diethyl ether, THF or dimethylformamide (DMF).

20 When G is chloro or bromo, the reaction of the compound of formula X with the appropriate compound of formula NH₂R³ is typically carried out in the presence of an acid scavenger in an aprotic solvent at a temperature from about 0°C to about 100°C. Suitable acid scavengers include triethylamine (TEA), pyridine and inorganic salts such as sodium and potassium carbonate. Suitable solvents include methylene chloride (CH₂Cl₂), chloroform (CHCl₃), benzene, toluene and tetrahydrofuran (THF). Preferably, the reaction is conducted in CH₂Cl₂ at room temperature using TEA as the acid scavenger.

-28-

When G is O-(C₁-C₆)alkyl, the reaction of the compound of formula NH₂R³ is usually conducted in an aprotic solvent such as benzene, toluene, chlorobenzene or xylenes, at a temperature from about 25°C to about 100°C, preferably at about the reflux temperature of the solvent.

Reduction of the compound of formula XII so formed yields the corresponding compound of the formula I wherein R² is hydrogen. This is generally accomplished using a reducing agent such as lithium aluminum hydride, borane dimethylsulfide complex, borane-THF or diborane, in an aprotic solvent such as THF, dioxane or diethyl ether, at a temperature from about 0°C to about 70°C. Preferably, the reducing agent is borane dimethylsulfide complex and the reaction is carried out at about room temperature in an ethereal solvent such as THF.

Compounds of the formula Ib may also be prepared by the procedures depicted in schemes 1 and 2 and described above, with the exception that ring system A in intermediate compounds X, XI and XII is replaced with the fused bicyclic ring system of the desired compound of formula Ib and R¹ is absent.

Compounds of the formula I wherein R² is hydrogen may be converted into the corresponding compounds wherein R² is -CO₂(C₁-C₁₀)alkyl by reacting them with a (C₁-C₁₀)alkyl halocarbonate such as methyl or ethyl chloroformate in the presence of an acid scavenger. Typically, this reaction is conducted in an polar solvent such as chloroform, methylene chloride, water or a water/acetone mixture, at a temperature from about 0°C to about 100°C, preferably at about room temperature. Suitable acid scavengers include triethylamine, pyridine and potassium and sodium carbonate or bicarbonate.

When R³ is a group of the formula II, the starting materials of the formula NH₂R³ may be prepared as described in United States Patent 5,162,339, which issued on November 11, 1992. This patent is incorporated herein in its entirety.

-29-

When R^3 is a group of the formula III, the starting materials of the formula NH_2R^3 may be prepared as described in United States Patent Application Serial No. 532,525, filed June 1, 1990 and PCT Patent Application PCT/US 5 91/02853, filed April 25, 1991. Both these applications are incorporated herein in their entirety.

When R^3 is a group of the formula IV, V or VI, the starting materials of the formula NH_2R^3 may be prepared as described in United States Patent Application Serial No. 10 557,442, filed July 23, 1990 and PCT Patent Application PCT/US 91/03369, filed May 14, 1991. Both these applications are incorporated herein in their entirety.

When R^3 is a group of the formula VII, the starting materials of the formula NH_2R^3 may be prepared as described 15 in United States Patent Application Serial No. 724,268, filed July 1, 1991, United States Patent Application Serial No. 800,667, filed November 27, 1991 and PCT Patent Application PCT/US 92/00065, filed January 14, 1992. These applications are incorporated herein in their entirety.

20 When R^3 is a group of the formula VIII, the starting materials of the formula NH_2R^3 may be prepared as described in PCT Patent Application PCT/US 91/05776, filed August 20, 1991, United States Patent Application Serial No. 800,667, filed November 27, 1991 and PCT Patent Application PCT/US 25 92/00065, filed January 14, 1992. These applications are incorporated herein in their entirety.

When R^3 is a group of the formula IX, the starting materials of the formula NH_2R^3 may be prepared as described in United States Patent Application Serial No. 719,884, 30 filed June 21, 1991. This application is incorporated herein in its entirety.

Scheme 2 illustrates one method of preparing the starting materials of formula X wherein G is hydrogen. This is the preferred method of preparing compounds of the 35 formula X wherein G is hydrogen and R^1 is thiazolyl, thiadiazolyl and oxazolyl. Once formed, these compounds can

-30-

be converted into the corresponding compounds of the formula I or XI according to the procedures described above.

Referring to scheme 2, a compound of the formula XIII is reacted with titanium tetrachloride (TiCl_4) or tin tetrachloride (SnCl_4) and dichloromethyl methyl ether ($\text{CHCl}_2\text{-O-CH}_3$) at a temperature from about 0°C to about room temperature, preferably at about 0°C , in a methylene chloride or tetrachloroethylene solvent to yield the corresponding aldehyde of formula X wherein G is hydrogen.

10 Alternatively, the compound of the formula XIII may be reacted with hexamethylene tetraamine and trifluoroacetic acid at a temperature from about 25°C to about 80°C , preferably at about 70°C , to yield the same product.

Scheme 3 illustrates a preferred method of preparing compounds of the formula X wherein G is hydrogen and R^1 is a nitrogen containing heterocyclic group (e.g., a pyrrolyl, triazolyl or imidazolyl group). Referring to scheme 3, the -CHO group of a benzaldehyde of the formula XIV is protected by conversion to the corresponding 1,3-dioxolane of formula

15 XV, wherein R^5 is a suitable leaving group such as iodine or bromine. This reaction is generally carried out by heating a mixture of the benzaldehyde and ethylene glycol in an inert solvent such as benzene or toluene, preferably in the presence of an acid catalyst such as p-toluenesulfonic acid,

20 and preferably at the reflux temperature of the solvent to remove the water formed in the reaction.

The resulting compound of formula XV is then reacted with a heterocyclic compound of the formula R^1H to form the corresponding compound of formula XVI. Typically, the

30 reaction is carried out in an aprotic, nonpolar solvent such as xylene or toluene, or in the absence of a solvent (e.g., as a melt of imidazole and the compound of the formula XV) at a temperature from about 100°C to about 300°C , in the presence of an inorganic metal catalyst such as copper metal


35 or copper iodide, in a high pressure reactor at a pressure from about 1 atm to about 5 atm. Preferably, the reaction is carried out neat using a copper metal catalyst, at a

-31-

temperature from about 140°C to about 160°C and at a pressure from about 2 atm to about 3 atm.

Treatment of the compound of formula XVI formed in the above reaction with a mixture of aqueous hydrochloric acid in acetone at a temperature from about 0°C to about 50°C, preferably at room temperature, will convert the dioxolane to the desired compound of formula X.


Alternatively, compounds of the formula X wherein G is hydrogen and R¹ does not contain an ionizable proton (for example, R¹ = 2-pyridyl, 2-thienyl or 2-pyrimidinyl) can be prepared by reacting a compound of the formula XV, as depicted in scheme 3 and defined above, wherein R⁵ is bromine or iodine, with magnesium metal to form a Grignard reagent of the formula



and then reacting the Grignard reagent in situ with a halogen substituted heterocyclic compound of the formula X''R¹ wherein X'' is chloro, bromo or iodo under standard Grignard conditions. These reactions are typically conducted in an ethereal solvent (e.g., diethyl ether or tetrahydrofuran), at a temperature from about 0°C to about 70°C. They are preferably conducted at the reflux temperature of the solvent in the presence of a catalyst (e.g., tetrakis (triphenylphosphine) palladium (0)).

Compounds of the formula X wherein G is other than hydrogen can be prepared from commercially available sources by methods well known to those skilled in the art. For example, compounds of the formula X wherein G is hydroxy can be obtained by: (1) oxidizing the corresponding compounds of the formula


-32-



5

XVII

wherein R^{20} is methyl with potassium permanganate in a
 10 reaction inert solvent such as acetone; (2) oxidizing an
 alcohol of the formula XVII wherein R^{20} is hydroxymethyl with
 manganese dioxide; or (3) subjecting a compound of the
 formula XVII wherein R^{20} is chloro, bromo or iodo to Grignard
 reaction conditions (i.e., reacting the compound of formula
 15 XVII with magnesium metal to form an intermediate of the
 formula



20

and then treating the intermediate with carbon dioxide.

The foregoing carboxylic acids of the formula X wherein
 25 G is hydroxy can be converted into the corresponding
 compounds of the formula X wherein G is chlorine or bromine
 by reaction with such reagents as sulfonyl chloride,
 phosphorus trichloride, phosphorus pentachloride and
 phosphorous tribromide.


30 Carboxylic esters of the formula X wherein G is (C_1-C_6) alkoxy can be prepared by a variety of methods known in
 the art. One such method involves reacting the
 corresponding acid halide in a (C_1-C_6) alkanol in the presence
 of a catalytic amount of hydrochloric, sulfuric or para-
 35 toluenesulfonic acid at a temperature from about room
 temperature to about the boiling point of the alcohol
 employed.

-33-

Compounds of the formula Ia wherein R¹ is pyrrolyl can also be prepared from the corresponding compounds wherein R¹ is replaced by an amino group. The corresponding amine may be obtained by reducing the corresponding nitro compound using one of several methods known to those skilled in the art. One such method involves catalytic hydrogenation of the nitro compound using hydrogen gas and a palladium on carbon catalyst in an inert solvent such as methanol or ethanol at about room temperature and a pressure of about 1-5 atm. The reduction can also be accomplished using a reducing agent such as borane/methyl sulfide in tetrahydrofuran at a temperature from about 25°C to about 70°C, preferably at the reflux temperature of the solvent. The latter reduction method is exemplified in Example 45 of United States Patent Application Serial No. 932,392, filed on August 19, 1992. This application is incorporated herein by reference in its entirety.

The amine can then be converted into the desired compound of formula I by the procedure described in Example 9.

Scheme 4 illustrates a method of preparing compounds of the formula



XXVIII

wherein R²² is methyl, ethyl, n-propyl, isopropyl, t-butyl, trifluoromethyl, (C₁-C₆)alkyl or benzyl, and R²³ is methyl, ethyl, propyl, isopropyl, t-butyl, trifluoromethyl, (C₁-C₆)alkyl, benzyl or phenyl optionally substituted with from one to three substituents independently selected from halo, trifluoromethoxy, (C₁-C₆)alkyl and (C₁-C₆)alkoxy. These

-34-

aldehydes are intermediates in the synthesis of compounds of the formula Ib wherein W is OR^{22} , X' is -S-, Y' is CR^{23} and Z' is =N-. Such compounds of the formula Ib may be prepared from the foregoing aldehydes of formula XVIII as described above and depicted in scheme 1.

Referring to scheme 4, compounds of the formula XX may be prepared by direct alkylation of the corresponding phenols of the formula XIX using a common alkylating agent such as dimethyl sulfate, a methyl halide (e.g., methyl iodide), methyl triflate, methyl mesylate or methyl tosylate. The reaction is usually conducted in an inert solvent such as dimethyl formamide, N-methyl pyrrolidinone, tetrahydrofuran, methylene chloride or another similar solvent for a period of about 0.5 to 12 hours at a temperature of about 0°C to the reflux temperature of the solvent. Typically, a base such as sodium hydride or potassium hydride is used, but other bases such as triethylamine, 1,8-diazobicyclo[5.4.0]undec-7-ene may be utilized as well. During the alkylation process, the carboxylic acid functionality of the phenol of formula XIX is also alkylated. However, various esterified derivatives of the compounds of formula XIX such as the methyl or ethyl ester (not shown in scheme 4) are also suitable starting materials for the foregoing transformation.

Reduction of the nitro functionality in the resulting compound of formula XX to yield the corresponding amine of formula XXI may be effected through hydrogenation with a noble metal catalyst such as platinum or palladium under a pressure of about 1-100 atmospheres of hydrogen gas in an inert solvent such as methanol, ethanol, ether, tetrahydrofuran or water (or a mixture of two or more such solvents). The reaction is most conveniently run at ambient temperature for about 0.5 to 12 hours. Alternatively, reduction of the nitro functionality may be carried out using a metal such as zinc or tin in a solvent such as acetic acid or water.

-35-

Formation of the amide of formula XXII is most conveniently conducted in an inert solvent such as methylene chloride, dichloroethane, tetrahydrofuran or toluene using one or more equivalents of an acylating agent such as acetic anhydride, acetyl chloride, benzoyl chloride, trimethylacetyl chloride, cyclopropyl carbonyl chloride or another alkyl or aryl acid chloride or anhydride. The reaction is most conveniently carried out in the presence of a base such as triethylamine or diisopropylamine or in aqueous solution under Schotten-Baumann conditions with sodium hydroxide. The reaction is generally conducted between about 0°C and 60°C, with room temperature being preferred.

The thioamide of formula XXIII is formed from the corresponding amide of formula XXII by reacting the latter compound with a reagent such as 2,4-bis(4-methoxyphenyl)-1,3-dithia-2,4-diphosphetane-2,4-disulfide (Lawesson's Reagent) or phosphorous pentasulfide in an inert solvent such as toluene, benzene, dichloroethane, dimethylformamide or hexamethylphosphorous triamide at a temperature from about room temperature to the reflux point of the solvent. The reaction may alternatively be run neat (without a solvent).

Cyclization of the thioamide of formula XXIII to form a mixture of regioisomeric benzothiazoles of the formulae XXIV and XXV is carried out by reacting the substrate with potassium ferricyanide in an aqueous base heated to about 50°C for a period of about 1 to 12 hours. The mixture of regioisomers may be separated at this point or carried through to the end of the synthetic sequence.

Reduction of the mixture of compounds of the formulae XXIV and XXV or of the compound of formula XXIV alone can be conducted using a reducing agent such as lithium aluminum hydride, borane-THF, sodium bismethoxyethylaluminum hydride or a similar reducing agent in an inert solvent such as ether, tetrahydrofuran, dimethoxyethane or toluene. The reaction may be carried out at a temperature between about

-36-

0°C and room temperature for a period of about 1 to 12 hours.

The resulting mixture of the regioisomeric alcohols of the formulae XXVI and XXVII (or XXVI separately) may be oxidized to form the desired aldehyde of formula XXVIII by reacting it with manganese dioxide in refluxing methylisobutyl ketone or another inert solvent for a period of about 1-12 hours. Alternatively, the oxidation can be carried out under "Swern" conditions (i.e., a mixture of dimethyl sulfoxide in methylene chloride with activation by oxalyl chloride or trifluoroacetic anhydride) or using other methods well known to those familiar with the art. Also effective are oxidizing agents such as pyridinium chlorochromate and pyridinium dichromate in an inert solvent such as methylene chloride, chloroform or dichloroethane.

The preparation of other compounds of the formula I not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

In each of the reactions discussed or illustrated in schemes 1 to 4 above, pressure is not critical unless otherwise indicated. Pressures from about 0.5 atmospheres to about 5 atmospheres are generally acceptable, and ambient pressure, i.e. about 1 atmosphere, is preferred as a matter of convenience.

The novel compounds of the formula I and the pharmaceutically acceptable salts thereof are useful as substance P antagonists, i.e., they possess the ability to antagonize the effects of substance P at its receptor site in mammals, and therefore they are able to function as therapeutic agents in the treatment of the aforementioned disorders and diseases in an afflicted mammal.

The compounds of the formula I which are basic in nature are capable of forming a wide variety of different salts with various inorganic and organic acids. Although such salts must be pharmaceutically acceptable for

-37-

administration to animals, it is often desirable in practice to initially isolate a compound of the Formula I from the reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained.

Those compounds of the formula I which are also acidic in nature, e.g., where R^6 or R^{10} is carboxyphenyl, are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include the alkali metal or alkaline-earth metal salts and particularly, the sodium and potassium salts. These salts are all prepared by conventional techniques. The chemical bases which are used as reagents to prepare the pharmaceutically acceptable base salts of this invention are those which form non-toxic base salts with the acidic compounds of formula I. Such non-toxic base salts include those derived from such pharmacologically acceptable cations as sodium, potassium, calcium and magnesium, etc. These salts can easily be prepared by treating the corresponding acidic compounds with an aqueous solution containing the desired pharmacologically acceptable cations, and then evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively, they may also be prepared by mixing lower alkanolic solutions of the acidic compounds and the desired alkali metal alkoxide together, and then evaporating the resulting solution to dryness in the same manner as before. In either case, stoichiometric quantities of reagents are

-38-

preferably employed in order to ensure completeness of reaction and maximum yields of the desired final product.

The compounds of formula I and their pharmaceutically acceptable salts exhibit substance P receptor-binding activity and therefore are of value in the treatment and prevention of a wide variety of clinical conditions the treatment or prevention of which are effected or facilitated by a decrease in substance P mediated neurotransmission. Such conditions include inflammatory diseases (e.g., arthritis, psoriasis, asthma and inflammatory bowel disease), anxiety, depression or dysthymic disorders, urinary incontinence, gastrointestinal disorders such as emesis and colitis, psychosis, pain, allergies such as eczema and rhinitis, chronic obstructive airways disease, hypersensitivity disorders such as poison ivy, vasospastic diseases such as angina, migraine and Reynaud's disease, fibrosing and collagen diseases such as scleroderma and eosinophilic fascioliasis, reflex sympathetic dystrophy such as shoulder/hand syndrome, addiction disorders such as alcoholism, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders such as Alzheimer's disease, AIDS related dementia, diabetic neuropathy and multiple sclerosis, disorders related to immune enhancement or suppression such as systemic lupus erythematosus, and rheumatic diseases such as fibrositis. Hence, these compounds are readily adapted to therapeutic use as substance P antagonists for the control and/or treatment of any of the aforesaid clinical conditions in mammals, including humans.

The compounds of the formula I and the pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in dosages ranging from about 5.0 mg up to about 1500 mg per day, although variations will necessarily occur depending upon the weight and condition of the subject being treated and the particular route of administration chosen. However, a

-39-

dosage level that is in the range of about 0.07 mg to about 21 mg per kg of body weight per day is most desirably employed. Variations may nevertheless occur depending upon the species of animal being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out. In some instances, dosage levels below the lower limit of the aforesaid range may be more than adequate, while in other cases still larger doses may be employed without causing any harmful side effect, provided that such larger doses are first divided into several small doses for administration throughout the day.

The compounds of the formula I and their pharmaceutically acceptable salts ("the therapeutic compounds") may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the three routes previously indicated, and such administration may be carried out in single or multiple doses. More particularly, the novel therapeutic agents of this invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutic compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine

-40-

may be employed along with various disintegrants such as starch (and preferably corn, potato or tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, 5 gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this 10 connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or 15 dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a 20 therapeutic compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous 25 injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well known to those skilled in the 30 art.

Additionally, it is also possible to administer the therapeutic compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, 35 pastes, ointments and the like, in accordance with standard pharmaceutical practice.

-41-

The activity of the therapeutic compounds of the present invention as substance P receptor antagonists may be determined by their ability to inhibit the binding of substance P at its receptor sites in bovine caudate tissue, employing radioactive ligands to visualize the tachykinin receptors by means of autoradiography. The substance P antagonizing activity of the herein described compounds may be evaluated by using the standard assay procedure described by M. A. Cascieri et al., as reported in the Journal of Biological Chemistry, Vol. 258, p. 5158 (1983). This method essentially involves determining the concentration of the individual compound required to reduce by 50% the amount of radiolabelled substance P ligands at their receptor sites in said isolated cow tissues, thereby affording characteristic IC₅₀ values for each compound tested.

In this procedure, bovine caudate tissue is removed from a -70°C freezer and homogenized in 50 volumes (w./v.) of an ice-cold 50 mM Tris (i.e., trimethamine which is 2-amino-2-hydroxymethyl-1,3-propanediol) hydrochloride buffer having a pH of 7.7. The homogenate is centrifuged at 30,000 x G for a period of 20 minutes. The pellet is resuspended in 50 volumes of Tris buffer, rehomogenized and then recentrifuged at 30,000 x G for another twenty-minute period. The pellet is then resuspended in 40 volumes of ice-cold 50 mM Tris buffer (pH 7.7) containing 2 mM of calcium chloride, 2 mM of magnesium chloride, 4 µg/ml of bacitracin, 4 µg/ml of leupeptin, 2 µg of chymostatin and 200 g/ml of bovine serum albumin. This step completes the production of the tissue preparation.

The radioligand binding procedure is then carried out in the following manner, viz., by initiating the reaction via the addition of 100 µl of the test compound made up to a concentration of 1 µM, followed by the addition of 100 µl of radioactive ligand made up to a final concentration 0.5 nM and then finally by the addition of 800 µl of the tissue preparation produced as described above. The final volume is thus 1.0 ml, and the reaction mixture is

-42-

next vortexed and incubated at room temperature (ca. 20°C) for a period of 20 minutes. The tubes are then filtered using a cell harvester, and the glass fiber filters (Whatman GF/B) are washed four times with 50 mM of Tris buffer (pH 7.7), with the filters having previously been presoaked for a period of two hours prior to the filtering procedure. Radioactivity is then determined in a Beta counter at 53% counting efficiency, and the IC₅₀ values are calculated by using standard statistical methods.

10 The ability of the therapeutic compounds of this invention to inhibit substance P induced effects in vivo may be determined by the following procedures "a" through "d". (Procedures "a" through "c" are described in Nagahisa et al., European Journal of Pharmacology, 217, 191-5 (1992), which is incorporated herein by reference in its entirety.)

15 a. Plasma extravasation in the skin

Plasma extravasation is induced by intradermal administration of substance P (50 µl, 0.01% BSA-saline solution) in dorsal skin of pentobarbital (25 mg/kg i.p.) anesthetized male Hartley guinea pigs weighing 450-500 g. The compound to be tested is dissolved in 0.1% methyl cellulose-water (MC) and dosed p.o. 1 hour before substance P challenge (3 pmol/site). Evans blue dye (30 mg/kg) is administered intravenously 5 minutes before challenge. After 10 minutes, the animals are sacrificed, the dorsal skin is removed, and the blue spots are punched out using a cork borer (11.5 mm oral dose (o.d.)). Tissue dye content is quantitated after overnight formamide extraction at 600 nm absorbance.

30 b. Capsaicin-induced plasma extravasation

Plasma extravasation is induced by intraperitoneal injection of capsaicin (10 ml of 30 µM solution in 0.1% BSA/saline) into pentobarbital anesthetized (25 mg/kg i.p.) guinea pigs. The compound to be tested is dissolved in 0.1% MC and dosed p.o. 1 hour before capsaicin challenge. Evans blue dye (30 mg/kg) is administered i.v. 5 minutes before challenge. After 10 minutes, the animals are sacrificed,

-43-

and both right and left ureters are removed. Tissue dye content is quantitated as in "a" above.

c. Acetic acid-induced abdominal stretching

Male ddY mice (SLC, Japan), weighing 14-18 g, were fasted overnight. The compound to be tested is dissolved in 0.1% MC and dosed p.o. 0.5 hour before acetic acid (AA) injection (0.7%, 0.16 ml/10 g body weight). The animals are placed in clear beakers (1 per beaker) and the stretching response is counted 10 to 20 minutes after the AA injection (10 minute interval).

d. Substance P-induced hyperlocomotor paradigm

The anti-psychotic activity of the therapeutic compounds of the present invention as neuroleptic agents for the control of various psychotic disorders may be determined by a study of their ability to suppress substance P-induced or substance P agonist induced hypermotility in guinea pigs. This study is carried out by first dosing the guinea pigs with a control compound or with an appropriate test compound of the present invention, then injecting the guinea pigs with substance P or a substance P agonist by intracerebral administration via canula and thereafter measuring their individual locomotor response to said stimulus.

The present invention is illustrated by the following examples. It will be understood, however, that the invention is not limited to the specific details of these examples.

PREPARATION 1

2-Methoxy-5-(1,2,3-thiadiazol-4-yl)benzaldehyde

A mixture of 0.99 grams (5.15 mmol) of 4-(4-methoxyphenyl)-1,2,3-thiadiazole (Maybridge Chemical Co.) in 23 mL of anhydrous methylene chloride (CH_2Cl_2) cooled to 0°C, was treated with 2.3 mL (21.9 mmol) of titanium tetrachloride and stirred for 30 min. The red solution was treated with 0.97 mL (10.7 mmol) of α,α -dichloromethyl methyl ether and allowed to warm to room temperature overnight. The reaction mixture was then poured over 100 mL of saturated aqueous sodium bicarbonate (NaHCO_3), the pH was adjusted to 7-8 with solid NaHCO_3 , and the solution was then

-44-

extracted with CH_2Cl_2 . The organic layer was dried with magnesium sulfate (MgSO_4) and concentrated to a yellow solid. Chromatography on silica gel (20% EtOAc: 80% Hexane) gave the pure title compound as a light yellow solid, 0.35 g
5 (31%).

M.P. 156-157 °C.

^1H NMR ($\text{DMSO}-d_6$) δ 4.0 (s, 3H), 7.4 (d, 1H), 8.4 (m, 2H), 9.7 (s, 1H), 10.4 (s, 1H).

In the same manner, the following aldehyde
10 intermediates were prepared:

2-Methoxy-5-(2-thiazolyl)benzaldehyde, 36%

M.P. 119-120°C.

^1H NMR (CDCl_3) δ 4.0 (s, 3H), 7.1 (d, 1H), 7.3 (d, 1H), 7.8 (d, 1H), 8.2 (dd, 1H), 8.3 (d, 1H), 10.5 (s, 1H).

15 2-Methoxy-5-(4-methyl-2-thiazolyl)benzaldehyde, 35%, oil

^1H NMR (CDCl_3) δ 2.46 (s, 3H), 3.95 (s, 3H), 6.82 (s, 1H), 7.02 (d, $J=8.7$ Hz, 1H), 8.15 (dd, 1H), 8.26 (d, $J=2.4$ Hz, 1H), 10.46 (s, 1H).

2-Methoxy-5-(5-oxazolyl)benzaldehyde, 22%

20 ^1H NMR (CDCl_3) δ 4.0 (s, 3H), 7.1 (d, 1H), 7.4 (s, 1H), 7.8 (dd, 1H), 7.9 (s, 1H), 8.1 (d, 1H), 10.5 (s, 1H).

2-Methoxy-5-(6-methylpyridin-2-yl)benzaldehyde

^1H NMR (CDCl_3 , free base) δ 2.6 (s, 3H), 3.95 (s, 3H), 7.0 (m, 2H), 7.4 (d, 1H), 7.6 (d, 1H), 8.3 (dd, 1H), 8.5 (d, 1H), 10.5 (s, 1H).
25

Mass Spectrum (m/e , %): 228 (M^+), 227 (M^+ , 40), 85 (100).

2-Methoxy-5-(pyridin-2-yl)benzaldehyde

^1H NMR (CDCl_3 , free base) δ 4.0 (s, 3H), 7.1 (d, 1H),
30 7.2 (q, 1H), 7.7 (d, 2H), 8.3 (dd, 1H), 8.4 (d, 1H), 8.7 (d, 1H), 10.5 (s, 1H).

2-Methoxy-5-(pyridin-3-yl)benzaldehyde

M.p. 77-79°C.

^1H NMR (CDCl_3 , free base) δ 4.0 (s, 3H), 7.2 (d, 1H),
35 7.4 (m, 1H), 7.8 (dd, 1H), 7.9 (dd, 1H), 8.1 (d, 1H), 8.6 (dd, 1H), 8.9 (d, 1H), 10.5 (s, 1H).

-45-

2-Methoxy-5-(pyrimidin-2-yl)benzaldehyde

¹H NMR (CDCl₃, free base) δ 4.0 (s, 3H), 7.1 (d, 1H), 7.2 (t, 1H), 8.7 (d, 1H), 8.8 (d, 2H), 9.0 (d, 1H), 10.5 (s, 1H).

5 Mass Spectrum (m/e, %): 215 (M⁺, 100), 214 (M+, 35).

2-Methoxy-5-(3,5-dimethylpyrazol-1-yl)benzaldehyde

¹H NMR (CDCl₃, free base) δ 2.4 (s, 6H), 4.0 (s, 3H), 6.0 (s, 1H), 7.05 (d, 1H), 7.6 (dd, 1H), 7.85 (d, 1H), 10.5 (s, 1H).

10 Mass Spectrum (m/e, %): 231 (M⁺, 100).

2-Methoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzaldehyde

M.p. 115-117°C.

¹H NMR (CDCl₃, free base) δ 1.9 (s, 3H), 2.1 (d, 6H), 4.0 (s, 3H), 7.0 (d, 1H), 7.8 (m, 1H), 7.9 (d, 1H), 10.5 (s, 1H).

15

Mass Spectrum (m/e, %): 245 (M⁺, 100).

2-Isopropoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzaldehyde

¹H NMR (CDCl₃, free base) δ 1.4 (d, 6H), 2.0 (s, 3H), 2.17 (s, 3H), 2.2 (s, 3H), 4.70 (m, 1H), 7.05 (d, 1H), 7.6 (dd, 1H), 7.8 (d, 1H), 10.5 (s, 1H).

20

2-Methoxy-5-(3,5-diisopropylpyrazol-1-yl)benzaldehyde

¹H NMR (CDCl₃, free base) δ 1.2 (d, 6H), 1.3 (d, 6H), 3.0 (m, 2H), 4.0 (s, 3H), 6.1 (s, 1H), 7.0 (d, 1H), 7.6 (dd, 1H), 7.9 (d, 1H), 10.5 (s, 1H).

25 Mass Spectrum (m/e, %): 286 (m+, 75), 271 (100).

5-(3,5-Dimethylthiophen-2-yl)-2-methoxybenzaldehyde

¹H NMR (CDCl₃, free base) δ 2.25 (s, 3H), 2.5 (s, 3H), 3.9 (s, 3H), 6.9 (d, 1H), 7.15 (d, 1H), 7.25 (m, 2H), 9.8 (s, 1H).

30 Mass Spectrum (m/e, %): 246 (M+, 100), 231 (35).

6-Methoxy-2,3-dimethyl-benzo[b]thiophene-7-carboxaldehyde

M.p. 155-160°.

¹H NMR (CDCl₃, free base) δ 2.2 (s, 3H), 2.45 (s, 3H), 4.0 (s, 3H), 7.0 (d, 1H), 7.8 (d, 1H), 10.7 (s, 1H).

35 Mass Spectrum (m/e, %): 221 (M⁺, 100).

-46-

6-Methoxy-2-methyl-benzoxazol-5-ylaldehyde

6-Methoxy-2-methyl-benzoxazole (1.06 grams, 6.5 mmol) was taken up in 65 mL of dry methylene chloride and cooled to 0°C. Titanium tetrachloride (10.11 grams, 53.3 mmol) was added via syringe and the deep red solution was stirred for 30 min. Dichloromethyl methyl ether (4.63 grams, 40.3 mmol) was added dropwise and the reaction mixture darkened to a brown color. The reaction mixture was allowed to warm to room temperature and was stirred for 18 hours. The mixture was quenched into ice and water. The slurry was basified with aqueous saturated bicarbonate and extracted with methylene chloride. The organic phase was washed with saturated aqueous brine and then dried and evaporated in vacuo. The residue was chromatographed on silica eluting over a gradient of 20% ethyl acetate/hexanes to 50% ethyl acetate/hexanes. In addition to starting material, a more polar material, the desired aldehyde, was obtained in 69 mg. Further elutions led to the isolation of a regioisomeric aldehyde. The desired material displayed the following spectral data.

¹H NMR (250 MHz, CDCl₃) δ 10.48 (s, 1H), 8.10 (s, 1H), 7.06 (s, 1H), 3.98 (s, 3H), 2.62 (s, 3H).

PREPARATION 26-Methoxy-2-methyl-benzothiazol-5-ylaldehyde

6-Methoxy-2-methyl-benzothiazole [3.34 grams (18.63 mmol)] was taken up in 35 mL of trifluoroacetic acid and treated with 2.62 grams (18.63 mmol) of hexamethylenetetramine. The reaction was heated under reflux for seven hours. The reaction mixture was allowed to cool and was evaporated in vacuo. The residue was diluted with 200 mL of ethyl acetate and treated with 100 mL of saturated sodium bicarbonate solution. The organic phase was washed with saturated brine solution and was dried and evaporated. The residue was chromatographed on silica gel (elution with 20% ethyl acetate in hexanes) to provide 3.04 grams of recovered starting material and 10 mg of the desired aldehyde.

-47-

¹H NMR (250 MHz, CDCl₃) δ 10.52 (s, 1H), 8.36 (s, 1H), 7.38 (s, 1H), 3.99 (s, 3H), 2.82 (s, 3H).

In a similar manner, the following aldehyde intermediate was prepared.

5 6-Methoxy-3-methyl-benzo[d]isoxazol-5-ylaldehyde

6-Methoxy-3-methyl-benzo[d]isoxazole (1.4 g, 8.58 mmol) was taken up in 200 mL of trifluoroacetic acid and treated with 1.20 grams (8.6 mmol) of hexamethylenetetramine. The reaction was heated under reflux for 24 hours. The reaction mixture was allowed to cool and was evaporated in vacuo. The residue was diluted with 200 mL of methylene chloride and treated with 200 mL of saturated sodium bicarbonate solution. The organic phase was washed with saturated brine solution and was dried and evaporated. The residue was chromatographed on silica gel (elution with 15% ethyl acetate in hexanes) to provide 122.6 mg of the desired aldehyde.

¹H NMR (250 MHz, CDCl₃) δ 10.46 (s, 1H), 8.15 (s, 1H), 7.04 (s, 1H), 4.02 (s, 3H), 2.56 (s, 3H).

20

PREPARATION 3

5-(2-Imidazolyl)-2-methoxybenzaldehyde

To a solution of 2.04 grams (8.72 mmol) of 4-iodoanisole in 36 mL of CH₂Cl₂, cooled to 0°C, was added dropwise 2.0 mL (18.7 mmol) of titanium tetrachloride. After stirring for 30 min., 0.93 mL (10.3 mmol) of α,α-dichloromethyl methyl ether was added and the reaction maintained at 0°C for another 2 hours. The reaction mixture was then poured with stirring into a mixture of 50 mL of CH₂Cl₂ and 50 mL of saturated aqueous sodium bicarbonate (NaHCO₃). After 30 min., this was filtered through diatomaceous earth, the organic layer was separated, the aqueous layer was twice extracted with CH₂Cl₂, and all of the organic layers were combined, dried with MgSO₄ and concentrated in vacuo. The crude residue was recrystallized from ethanol (EtOH) to give 5-iodo-2-methoxybenzaldehyde as pale yellow needles, 1.37 grams (60%).

-48-

¹H NMR (CDCl₃) δ 4.0 (s, 3H), 6.8 (d, 1H), 7.8 (dd, 1H), 8.1 (d, 1H), 10.4 (s, 1H).

A solution of 1.0 grams (3.82 mmol) of the above aldehyde, 0.85 mL of ethylene glycol, 20 mg of p-toluenesulfonic acid and 42 mL of toluene was refluxed under nitrogen for 18 hours, cooled and concentrated in vacuo. The residue was redissolved in 50 mL of CH₂Cl₂, washed with saturated aqueous NaHCO₃, dried over MgSO₄ and concentrated to an oil. Chromatography on silica gel (10% ethyl acetate (EtOAc): 90% hexane) gave 2-(5-iodo-2-methoxyphenyl)-1,3-dioxolane as a clear oil, 0.68 grams (58%).

¹H NMR (CDCl₃) δ 3.9 (s, 3H), 4.0-4.3 (m, 4H), 6.1 (s, 1H), 6.7 (d, 1H), 7.4 (dd, 1H), 7.9 (d, 1H).

In a Pyrex high pressure reaction tube, a solution of the preceding dioxolane (200 mg, 0.66 mmol), 140 mg (2.06 mmol) of imidazole and 90 mg (1.4 mmol) of copper powder in 1 mL of anhydrous tetrahydrofuran (THF) was stirred until homogeneous. The solvent was then evaporated under nitrogen, and the tube was sealed and heated in an oil bath at 140°C for 16 hours. After cooling, the residue was dissolved in 20 mL of THF, filtered through a pad of diatomaceous earth and concentrated in vacuo to an oil. Chromatography on silica gel (100% ethyl acetate (EtOAc)) gave 60 mg of 2-(5-(2-imidazolyl)-2-methoxyphenyl)-1,3-dioxolane as a yellow oil.

MS: m/e 246 (m+).

The preceding oil in 20 mL of acetone was treated with 10 mL of 1 N hydrochloric acid (HCl) and stirred at 25°C for 3 hours, evaporated in vacuo and extracted into CH₂Cl₂. After washing with water, the organic layer was dried over MgSO₄ and concentrated to a clear oil. Chromatography on silica gel (3% CH₃OH: 97% CH₂Cl₂) gave 5-(2-imidazolyl)-2-methoxybenzaldehyde as a clear oil, 30 mg (61%).

¹H NMR (CDCl₃) δ 4.0 (s, 3H), 7.1 (d, 1H), 7.1-7.3 (m, 2H), 7.6 (dd, 1H), 7.8 (d, 2H), 10.5 (s, 1H).

-49-

EXAMPLE 1(2S,3S-3-[5-(2-Imidazolyl)-2-methoxybenzyl]amino-2-phenylpiperidine trihydrochloride dihydrate

Under a nitrogen atmosphere, a mixture of 30 mg (0.15 mmol) of 5-(2-imidazolyl)-2-methoxybenzaldehyde and 54 mg (0.15 mmol) of (+)-(2S,3S)-3-amino-2-phenylpiperidine in 5 mL of dry toluene was stirred and heated to reflux for 18 hours, using a Dean-Stark trap. The toluene in the reaction flask was removed in vacuo, the residue was dissolved in 5 mL of anhydrous 1,2-dichloroethane and stirred for 10 min. Sodium triacetoxyborohydride (92 mg, 0.43 mmol) was added and stirring was continued overnight. The solvent was removed in vacuo and the residue was treated with 5 mL of water (H₂O) and extracted with methylene chloride (CH₂Cl₂) (3 x 10 mL). The organic extracts were combined, dried over magnesium sulfate (MgSO₄) and concentrated to an oil. Chromatography on silica gel using methylene chloride (CH₂Cl₂): Methanol (CH₃OH): concentrated ammonium hydroxide (NH₄OH) in a ratio of 94:5:1 gave the pure free base as a clear oil. The oil dissolved in CH₂Cl₂ was treated with a solution of Et₂O saturated with hydrogen chloride (HCl) gas to give the title product as a pale yellow solid, 15.7 mg (21%).

M.P. 235°C (decomp.)

¹H NMR (CDCl₃, free base) δ 1.5 (m, 1H), 1.7 (m, 1H), 1.9 (m, 4H), 2.2 (m, 1H), 2.9 (m, 2H), 3.3 (m, 1H), 3.5 (d, 1H), 3.6 (s, 3H), 3.7 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.0 (d, 1H), 7.1-7.4 (m, 8H), 7.7 (s, 1H).

FAB MS: m/e 363 (m+1)

Anal. calc'd for C₂₂H₂₆N₄O•3HCl•2H₂O: C, 52.03; H, 6.55; N, 11.03. Found: C, 51.84; H, 6.36; N, 10.53.

The title compounds of Examples 2 through 15 were prepared from (+)-(2S,3S)-3-amino-2-phenylpiperidine and the appropriate aldehyde using a procedure similar to that of Example 1.

-50-

EXAMPLE 2(2S,3S)-3-[2-Methoxy-5-(2-thiazolyl)benzyl]amino-2-phenylpiperidine trihydrochloride dihydrate

M.P. 207°C (decomp.)

5 ¹H NMR (CDCl₃, free base) δ 1.5 (m, 1H), 1.7 (m, 1H), 2.0 (m, 4H), 2.8 (m, 1H), 2.8 (m, 2H), 3.3 (m, 1H), 3.4 (d, 1H), 3.5 (s, 3H), 3.8 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.3 (m, 7H), 7.6 (d, 1H), 7.8 (m, 2H).

FAB MS: m/e 380 (m+1)

10 Anal. calc'd for C₂₂H₂₅N₃OS•3HCl•2H₂O: C, 50.34; H, 6.14; N, 8.00. Found: C, 50.24; H, 6.05; N, 7.84.

EXAMPLE 3(2S,3S)-3-[2-Methoxy-5-(1,2,3-thiadiazol-4-yl)benzyl]amino-2-phenylpiperidine trihydrochloride dihydrate

15 M.P. > 270°C (decomp.)

¹H NMR (CDCl₃, free base) δ 1.4 (m, 1H), 1.6 (m, 1H), 1.9 (m, 3H), 2.2 (m, 2H), 3.2 (m, 2H), 3.5 (d, 4H), 3.7 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.3 (m, 5H), 7.6 (d, 1H), 7.9 (dd, 1H), 8.4 (s, 1H).

20 FAB MS: m/e 381 (m+1, 100%), 353, 321.

EXAMPLE 4(2S,3S)-3-[2-methoxy-5-(5-oxazolyl)benzyl]amino-2-phenylpiperidine dihydrochloride

M.P. 267°C (decomp.)

25 ¹H NMR (CDCl₃, free base) δ 1.5 (m, 1H), 1.7 (m, 1H), 2.0 (m, 4H), 2.2 (m, 1H), 2.9 (m, 2H), 3.3 (m, 1H), 3.5 (d, 1H), 3.6 (s, 3H), 3.7 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.2 (s, 1H), 7.3 (m, 6H), 7.5 (dd, 1H), 7.9 (s, 1H).

FAB MS: m/e 364 (m+1)

30 Anal. calc'd for C₂₂H₂₅N₃O₂•2HCl: C, 60.55; H, 6.24; N, 9.63. Found: C, 60.48; H, 6.21; N, 9.86.

EXAMPLE 5(2S,3S)-3-[2-Methoxy-5-(4-methyl-2-thiazolyl)benzyl]amino-2-phenylpiperidine trihydrochloride

35 M.P. 239°C (decomp.)

¹H NMR (CDCl₃, free base) δ 1.5 (m, 1H), 1.7 (m, 1H), 2.2 (m, 1H), 2.6 (s, 3H), 2.9 (m, 2H), 3.3 (m, 1H), 3.4 (d,

-51-

1H), 3.5 (s, 3H), 3.6 (d, 1H), 3.8 (d, 1H), 6.7 (d, 1H), 6.8 (s, 1H), 7.2-7.4 (m, 5H), 7.6 (d, 1H), 7.8 (dd, 1H).

FAB MS: m/e 394 (m+1).

EXAMPLE 6

5 (2S,3S)-(2-Methoxy-5-pyridin-2-ylbenzyl)-(2-phenylpiperidin-3-yl)amine trihydrochloride hydrate

M.p. 230°C.

¹H NMR (CDCl₃, free base) δ 1.35-2.2 (m, 6H), 2.8 (m, 2H), 3.3 (m, 1H), 3.5 (s, 3H), 3.6 (d, 1H), 3.8 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.0-7.3 (m, 6H), 7.5 (d, 1H), 7.6 (m, 2H), 7.8 (dd, 1H), 8.6 (M+1, 100).

Mass Spectrum (m/e, %): 374 (M+1, 100).

Anal. calc'd for C₂₄H₂₇N₃O•3HCl•1/4H₂O: C, 58.69; H, 6.21; N, 8.56. Found: C, 58.60; H, 6.18; N, 8.37.

15

EXAMPLE 7

(2S,3S)-(2-Methoxy-5-pyrimidin-2-ylbenzyl)-(2-phenylpiperidin-3-yl)amine dihydrochloride hydrate

M.p. 207°C (dec.)

¹H NMR (CDCl₃, free base) δ 1.5-2.0 (m, 5H), 2.4 (m, 1H), 2.8 (m, 1H), 2.8 (m, 2H), 3.4 (m, 1H), 3.5 (s, 3H), 3.6 (d, 1H), 3.9 (d, 1H), 4.0 (d, 1H), 5.7 (d, 1H), 7.1 (t, 1H), 7.4 (m, 5H), 8.1 (d, 1H), 8.3 (dd, 1H), 8.7 (d, 2H).

Mass Spectrum (m/e, %): 375 (M+1, 100).

Anal. calc'd for C₂₃H₂₆N₄O•2HCl•2H₂O: C, 57.14; H, 6.67; N, 11.59. Found: C, 57.23; H, 6.50; N, 11.44.

25

EXAMPLE 8

(2S,3S)-(2-Methoxy-5-pyridin-3-ylbenzyl)-(2-phenylpiperidin-3-yl)amine trihydrochloride hydrate

M.p. 241°C (dec.)

¹H NMR (CDCl₃, free base) δ 1.35-2.20 (m, 6H), 2.8 (m, 2H), 3.3 (d, 1H), 3.5 (d, 1H), 3.55 (s, 3H), 3.75 (d, 1H), 3.9 (d, 1H), 6.8 (d, 1H), 7.15-7.40 (m, 8H), 7.75 (m, 1H), 8.5 (dd, 1H), 8.72 (d, 1H).

Mass Spectrum (m/e, %): 374 (M+1, 100).

Anal. calc'd for C₂₄H₂₇N₃O•3HCl•H₂O: C, 57.55; H, 6.44; N, 8.39. Found: C, 57.88; H, 6.67; N, 8.07.

35

-52-

EXAMPLE 9

(2S,3S)-[2-Methoxy-5-(6-methylpyridin-2-yl)benzyl]-(2-phenylpiperidin-3-yl)amine dihydrochloride hydrate

M.p. 215-220°C.

5 ¹H NMR (CDCl₃, free base) δ 1.35-2.25 (m, 6H), 2.65 (s, 3H), 2.85 (m, 2H), 3.3 (d, 1H), 3.5 (s, 3H), 3.55 (d, 1H), 3.8 (d, 1H), 3.95 (d, 1H), 6.8 (d, 1H), 7.00 (d, 1H), 7.3 (m, 5H), 7.45 (d, 1H), 7.6 (m, 2H), 7.9 (dd, 1H).

Mass Spectrum (m/e, %): 387 (M⁺, 5), 268, 213 (100).

10 Anal. calc'd for C₂₅H₂₉N₃O•2HCl•1/4H₂O: C, 65.58; H, 6.83; N, 9.04. Found: C, 64.80; H, 6.76; N, 8.95.

EXAMPLE 10

(2S,3S)-[5-(3,5-Dimethylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine dihydrochloride hydrate

15 M.p. 255-259°C.

¹H NMR (CDCl₃, free base) δ 1.4-2.2 (m, 5H), 2.15 (d, 1H), 2.2 (s, 3H), 2.3 (s, 3H), 2.8 (m, 2H), 3.25 (d, 1H), 3.45 (d, 1H), 3.5 (s, 3H), 3.75 (d, 1H), 3.9 (d, 1H), 6.0 (s, 1H), 6.7 (d, 1H), 7.05 (d, 1H), 7.15-7.35 (m, 6H).

20 Mass Spectrum (m/e, %): 391 (M⁺, 100), 232 (20).

Anal. calc'd for C₂₄H₃₀N₄O•2HCl•1/4H₂O: C, 61.60; H, 7.00; N, 11.97. Found: C, 61.60; H, 6.97; N, 11.72.

EXAMPLE 11

25 (2S,3S)-[2-Methoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine trihydrochloride

M.p. 220-225°C.

30 ¹H NMR (CDCl₃, free base) δ 1.4 (d, 1H), 1.55 (tt, 1H), 1.95 (m, 3H), 2.05 (s, 3H), 2.15 (s+m, 4H), 2.25 (s, 3H), 2.8 (m, 2H), 3.3 (d, 1H), 3.45 (d, 1H), 3.5 (s, 3H), 3.75 (d, 1H), 3.9 (d, 1H), 6.7 (d, 1H), 7.0 (d, 1H), 7.15-7.35 (m, 6H).

Mass Spectrum (m/e, %): 406 (M⁺, 100).

Anal. calc'd for C₂₅H₃₂N₄O•3HCl•1/2CH₂Cl₂: C, 55.04; H, 6.52; N, 10.07. Found: C, 54.80; H, 6.82; N, 9.74.

35

EXAMPLE 12

-53-

(2S,3S)-[2-Isopropoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine hydrochloride hydrate

M.p. 140-155°C.

5 ¹H NMR (CDCl₃, free base) δ 1.05 (dd, 6H), 1.45-2.0 (m, 6H), 2.05 (s, 3H), 2.15 (s, 3H), 2.25 (s, 3H), 2.80 (t, 1H), 2.9 (d, 1H), 3.3 (d, 1H), 3.4 (d, 1H), 3.65 (d, 1H), 3.9 (d, 1H), 4.4 (m, 1H), 6.75 (d, 1H), 7.0 (d, 1H), 7.1-7.35 (m, 6H).

10 Mass Spectrum (m/e, %): 433 (M+1, 100).

Anal. calc'd for C₂₇H₃₆N₄O•HCl•H₂O: C, 66.58; H, 8.07; N, 11.50. Found: C, 66.54; H, 8.05; N, 11.83.

EXAMPLE 13

15 (2S,3S)-[5-(3,5-Diisopropylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine dihydrochloride hydrate

M.p. 222°C dec.

¹H NMR (CDCl₃, free base) δ 1.15 (dd, 6H), 1.3 (d, 6H), 1.35-2.15 (m, 6H), 2.85 (m, 3H), 3.05 (m, 1H), 3.25 (d, 1H), 20 3.45 (d, 1H), 3.5 (s, 3H), 3.7 (d, 1H), 3.9 (d, 1H), 6.0 (s, 1H), 6.7 (d, 1H), 7.0 (d, 1H), 7.15-7.35 (m, 6H).

Mass Spectrum (m/e, %): 446 (M+, 40), 232 (100).

EXAMPLE 14

25 (2S,3S)-[5-(3,5-Dimethylthiophen-2-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine dihydrochloride

M.p. 252-254°C.

¹H NMR (CDCl₃, free base) δ 1.37-2.01 (m, 6H), 2.20 (s, 3H), 2.38 (d, 3H), 2.8 (m, 2H), 3.27 (d, 2H), 3.50 (d, 1H), 3.85 (s, 3H), 3.90 (d, 1H), 6.3 (d, 1H), 6.72 (d, 1H), 7.03 30 (m, 2H), 7.2-7.32 (m, 5H).

Mass Spectrum (m/e, %): 408 (M+2, 30), 407 (M+1, 100).

Anal. calc'd for C₂₅H₃₀N₂OS•2HCl: C, 62.62; H, 6.73; N, 5.84. Found: C, 62.29; H, 6.44; N, 5.91.

-54-

EXAMPLE 15

(2S,3S)-(6-Methoxy-2,3-dimethyl-benzo[b]thiophen-7-ylmethyl)-(2-phenylpiperidin-3-yl)amine dihydrochloride hydrate

5 M.p. 240-244°C.

¹H NMR (CDCl₃, free base) δ 1.35-2.10 (m, 5H), 2.25 (s, 3H), 2.3 (m, 1H), 2.4 (s, 3H), 2.8 (m, 2H), 3.3 (d, 1H), 3.5 (s, 3H), 3.75 (d, 1H), 3.9 (s+d, 2H), 6.8 (d, 1H), 7.2 (m, 5H), 7.35 (d, 1H).

10 Mass Spectrum (m/e, %): 381 (M+1, 100).

Anal. calc'd for C₂₃H₂₈N₂OS•HCl•1/4H₂O: C, 65.54; H, 7.05; N, 6.65. Found: C, 65.47; H, 7.02; N, 6.56.

EXAMPLE 16

(1SR,2SR,3SR,4RS)-(2-Benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-methyl-benzoxazol-5-ylmethyl)-amine

A. (1SR,2SR,3SR,4RS)-(2-Benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-methyl-benzoxazol-5-ylmethylene)-amine.

(1SR, 2SR, 3SR, 4RS)-(2-Benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-amine (54 mg, 0.194 mmol) was dissolved in toluene (35 mL) and was treated with 37 mg (0.194 mmol) of 6-methoxy-2-methylbenzoxazol-5-ylcarboxaldehyde. The reaction mixture was heated under reflux over a Dean-Stark trap for 26 hours. Analysis of the NMR spectrum from a small reaction aliquot indicated product formation was complete. The solution was evaporated in vacuo to provide the imine as a crude oil which was used directly in the next step without purification.

B. (1SR,2SR,3SR,4RS)-(2-Benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-methyl-benzoxazol-5-ylmethyl)-amine

The crude imine from above was taken into 25 mL of dichloroethane and treated with 58 mg (0.272 mmol) of sodium triacetoxyborohydride. The mixture was stirred overnight (16 hours). Thin layer analysis (CH₂Cl₂:methanol(MeOH): concentrated ammonium hydroxide(NH₄OH) - 94:5:1) indicated the reaction was complete. Reaction quenching with 20 mL of

-55-

saturated aqueous sodium bicarbonate solution was followed by dilution with methylene chloride, extraction and drying. The organic phase was evaporated in vacuo to afford 130 mg of an oil. The crude product was chromatographed on silica gel eluting with a mixture of methylene chloride, methanol and aqueous ammonium hydroxide (98:1:1) to afford 90 mg of free base. The dihydrochloride salt was formed after dissolution of the free base in ether and treatment with saturated HCl gas, also in ether. The crude salt was obtained by direct evaporation of this reaction mixture. The residue was taken up in methanol (1 mL), clarified and treated with ether until the cloud point. The mixture was stirred overnight, whereupon crystallization occurred. The resulting solid (35 mg) was isolated by filtration.

¹H NMR (250 MHz, CDCl₃) δ 7.35 - 7.06 (m, 12H), 6.81 (s, 1H), 4.18 (d, 1H, J=12.2 Hz), 3.76 (d, 1H, J=13.7Hz), 3.52 (s, 3H), 3.51-3.40 (obsc. m, 1H), 3.43 (d, 1H, J=13.5Hz), 3.08 (br.d, 1H, J=9.1Hz), 2.76-2.60 (obsc. m, 2H), 2.60 (s, 3H), 2.50-2.40 (m, 1H), 2.17 (d, 1H, J=9.5Hz), 1.75-1.55 (m, 2H), 1.12-1.0 (m, 1H).

¹³C NMR (62.9 MHz, CDCl₃) δ 162.4, 155.5, 150.7, 145.8, 143.7, 134.1, 129.0, 128.9, 128.5, 127.7, 127.3, 126.4, 125.8, 124.9, 119.6, 92.7, 72.1, 63.3, 56.1, 55.7, 54.9, 50.8, 47.8, 41.4, 26.9, 14.5.

IR dihydrochloride salt (neat) λ 3000 (br), 2550 (br), 1620, 1580, 1500, 1480, 1460, 1440, 1380, 1350, 1310, 1210, 1150, 1120 cm⁻¹.

FAB MS: m/e 454 (m+1).

The title compounds of Examples 17 and 18 were prepared using procedures analogous to those of Examples 1 through 16.

EXAMPLE 17

(1SR,2SR,3SR,4RS)-(2-Benzhydryl-1-aza-bicyclo[2.2.1]hept-3-yl)-(6-methoxy-3-methyl-benzo[d]isoxazol-5-ylmethyl)-amine

¹H NMR (250 MHz, CDCl₃) δ 7.30-7.06 (m, 12H), 7.00 (s, 1H), 6.81 (s, 1H), 4.28 (d, 1H, J=12.1Hz), 3.75 (d, 1H,

-56-

J=13.7Hz), 3.61 (s, 3H), 3.53 - 3.46 (obsc. m, 1H), 3.45 (d, 1H, J=13.7Hz), 3.07 (br.d, 1H, J=9.6Hz), 2.73 (dd, 2H, J=15.1 Hz, J=8.3Hz), 2.65 (d, 1H, J=4.9Hz), 2.53 (s, 3H), 2.53-2.41 (m, 1H), 2.18 (d, 1H, J=9.6Hz), 1.7-1.5 (m, 2H),
5 1.14-1.07 (m, 1H).

¹³C NMR (75.5 MHz, CDCl₃) δ 163.8, 159.9, 154.5, 145.5, 143.7, 128.8, 128.4, 127.7, 127.2, 126.8, 125.2, 125.5, 120.4, 114.5, 90.9, 72.0, 63.8, 55.9, 55.7, 54.7, 50.7, 47.7, 41.4, 26.8, 10.0.

10 IR (neat) λ 3340, 2950, 1620, 1600, 1490, 1470, 1450, 1380, 1340, 1270, 1190, 1150 cm⁻¹.

HRMS calc'd for C₂₉H₃₂N₃O₂: 454.2487. Found: 454.2522.

EXAMPLE 18

(2S,3S)-(6-Methoxy-2-methyl-benzothiazol-5-ylmethyl)-

15 (2-phenylpiperidin-3-yl)-amine

¹H NMR (250 MHz, D₂O) dihydrochloride salt δ 7.63 (s, 1H), 7.57-7.55 (m, 1H), 7.49-7.40 (m, 1H), 7.36-7.25 (m, 3H), 7.12-7.09 (m, 2H), 4.89-4.87 (d, 1H, J=3.6Hz), 4.47-4.42 (d, 1H, J=13.3Hz), 4.17-4.12 (d, 1H, J=13.2Hz), 4.05-4.02 (m, 20 1H), 3.71-3.61 (obsc. m, 1H), 3.61 (3H, s), 3.38-3.23 (m, 1H), 2.80 (3H, s), 2.54-2.47 (1H, m), 2.29-2.20 (2H, m), 2.11-1.99 (3H, m).

IR dihydrochloride salt (neat) λ 2700 br, 1610, 1560, 1520, 1460, 1430, 1420, 1280, 1220, 1050 cm⁻¹.

25 HRMS calc'd for C₂₁H₂₆N₃OS: 368.1791. Found: 368.17668.

-57-

EXAMPLE 19(2S,3S)-[5-(2,5-Dimethylpyrrol-1-yl)-2-methoxybenzyl]-
(2-phenylpiperidin-3-yl)amineA. 2-Methoxy-5-nitrobenzaldehyde

5 A mixture of 2.5 grams (13.5 mmol) of 2-chloro-5-nitrobenzaldehyde (Aldrich Chem. Co.) in 250 mL of CH₃OH was treated with 2.92 grams (54.1 mmol) of sodium methoxide, refluxed for 16 hours and allowed to cool to room temperature. After removal of the solvent in vacuo, the
10 residue was suspended in water, treated with 2N HCl (adjusting to pH < 5) and extracted into methylene chloride. After drying over MgSO₄, the solvent was removed in vacuo to give a yellow solid, 2.02 grams (83%).

¹H-NMR (CDCl₃) δ 4.1 (s, 3H), 7.2 (d, 1H), 8.5 (dd, 1H),
15 8.7 (d, 1H), 10.5 (s, 1H).

B. (2S,3S)-3-(2-Methoxy-5-nitrobenzyl)amino-2-phenyl-
piperidine

A mixture of 0.63 grams (3.48 mmol) of the above aldehyde and 0.60 grams (3.4 mmol) of (2S,3S)-3-amino-2-
20 phenylpiperidine (prepared as described in U.S. Patent application 724,268, filed on July 1, 1991) in 60 mL of toluene was refluxed under nitrogen for 18 hours with a Dean-Stark apparatus to remove the water generated in the reaction. The toluene was then removed in vacuo and the
25 oily residue was redissolved in 50 mL of methanol (CH₃OH), stirred under nitrogen for 1 hour and treated with 0.16 grams (4.23 mmol) of sodium borohydride. After stirring for 24 hours at 25°C, the solvent was evaporated in vacuo and the residue was treated with 50 mL of water, stirred for 1
30 hour and extracted with methylene chloride. The organic extracts were combined, dried over MgSO₄ and concentrated in vacuo to a pale orange oil. Chromatography on silica gel using concentrated ammonium hydroxide (NH₄OH): CH₃OH: CH₂Cl₂ (1:5:94) gave the title intermediate as a pale orange oil,
35 0.99 grams (85%).

¹H NMR (CDCl₃) δ 1.4 (d, 1H), 1.6 (tt, 1H), 1.7-1.95 (m, 3H), 2.05 (d, 1H), 2.75 (t, 2H), 3.15 (d, 1H), 3.35 (d, 1H),

-58-

3.55 (s, 1H), 3.65 (d, 1H), 3.85 (d, 1H), 6.65 (d, 1H), 7.2 (m, 5H), 7.9 (d, 1H), 8.0 (dd, 1H).

Mass Spectrum (m/e, %): 342 (m+1, 100%), 312 (10), 177 (16), 158 (14).

5 C. (2S,3S)-3-(5-Amino-2-methoxybenzyl)amino-2-phenyl-piperidine

A solution of 0.3 grams (0.88 mmol) of the preceding compound in 10 mL of ethyl acetate (EtOAc) was treated with 0.11 grams of 10% palladium on carbon and hydrogenated in a
10 Parr Shaker apparatus at an initial hydrogen pressure of 45 psi for a total of 4 hours. The reaction mixture was filtered through a pad of diatomaceous earth (d.e.) and the d.e. cake washed with additional EtOAc. The combined organics were concentrated in vacuo to give a pale orange
15 oil, 0.33 grams.

¹H NMR (CDCl₃) δ 1.5 (m, 1H), 1.7 (m, 1H), 2.0 (m, 1H), 2.2 (m, 1H), 2.7-2.9 (m, 6H), 3.3 (m, 1H), 3.4 (d+s, 4H), 3.6 (d, 1H), 3.9 (d, 1H), 6.3 (d, 1H), 6.4-6.6 (m, 2H), 7.2-7.4 (m, 5H).

20 D. (2S,3S)-[5-(2,5-Dimethylpyrrol-1-yl)-2-methoxybenzyl]-2-(2-phenylpiperidin-3-yl)amine

The above amine (0.33 grams) from part C in 12 mL of toluene was treated with 0.15 mL (1.28 mmol) of acetylacetone and refluxed with a Dean-Stark apparatus for
25 2 hours. After cooling to 25°C, the solvent was removed in vacuo to give a yellow oil. Chromatography on silica gel using concentrated NH₄OH: CH₃OH: CH₂Cl₂ (1:5:94) gave a clear oil, 198 mg. The oil was converted to the hydrochloride salt of the title compound by suspending the free base in
30 CH₂Cl₂ and treating it with 257 μL of 4M HCl in dioxane (2 equivalents), stirring at 25°C for 20 min, filtering and drying the solid in vacuo to give a light yellow solid, 0.19 grams.

M.P. 217°C (decomp).

35 ¹H NMR (CDCl₃, free base) δ 1.4 (m, 1H), 1.8 (m, 1H), 1.8 (bs, 1H), 1.9 (m, 1H), 2.0 (s, 6H), 2.1 (m, 1H), 2.8 (m, 2H), 3.3 (m, 1H), 3.4 (d, 1H), 3.6 (s, 3), 3.8 (d, 1H), 3.9

-59-

(d, 1H), 5.9 (s, 2H), 6.7 (d, 1H), 6.9 (d, 1H), 7.0 (dd, 1H), 7.3 (m, 5H).

Mass spectrum: (m/e, %) 390 (m+1), 270.

Anal. calc'd for $C_{25}H_{31}N_3O \cdot 2HCl \cdot 1/3CH_2Cl_2$: C, 62.00; H, 6.91; N, 8.56. Found: C, 62.07; H, 6.86; N, 8.21.

EXAMPLE 20

(2S, 3S)-(6-Methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenyl-piperidin-3-yl)-amine dihydrochloride salt

A. Methyl-2-methoxy-5-nitrobenzoate

10 A 500 ml round bottom flask was charged with 5.35 grams of 60% sodium hydride dispersion and the solid was washed twice with hexane and decanted (under nitrogen). The residue was suspended in 200 mL of dimethylformamide (DMF), cooled to 0°C and was treated with 11 grams (60 mmol) of 2-
15 hydroxy-5-nitrobenzoic acid. The reaction mixture was stirred for 1 hour at 0°C. The mixture was then treated with 17 mL (180 mmol) dimethyl sulfate and was allowed to warm to room temperature while stirring for 1.5 hours. An additional 17 mL of dimethyl sulfate was added and the
20 reaction was stirred overnight at room temperature. The mixture was diluted with 1.5 L ethyl acetate and was washed with 400 ml of 0.75 M aqueous sodium hydroxide (NaOH) solution followed by 400 mL of saturated bicarbonate solution, 400 mL of water and 500 mL of saturated brine.
25 The organic phase was dried and evaporated. There was obtained 11.6 grams of the desired product as yellow flakes. The material was generally used without purification.

Mass Spectrum: (m/e) 212 (m+1)

B. Methyl-2-methoxy-5-aminobenzoate

30 A 2 liter Parr Bottle was charged with 11.6 grams (55 mmol) methyl-2-methoxy-5-nitrobenzoate, 450 mL of methanol and 150 mL of tetrahydrofuran (THF). Platinum oxide was added (460 mg) and the mixture was hydrogenated under 45 psi hydrogen for 1 hour. The reaction mixture was filtered
35 through Celite® (diatomaceous earth) and evaporated. The reaction mixture was diluted with 500 mL of methylene chloride, and then washed with water and brine. The organic

-60-

phase was dried and evaporated (9.9 grams). This material was used without further purification.

Mass Spectrum m/e 181 (m+)

C. Methyl-2-methoxy-5-benzamidobenzoate

5 A solution of 9.92 grams (55 mmol) of methyl-2-methoxy-5-aminobenzoate was taken up in 250 ml methylene chloride, treated with 8.4 mL (61 mmol) triethylamine and cooled to 0°C. The reaction mixture was then treated with 8.1 grams (57 mmol) benzoyl chloride in a dropwise fashion. The
10 reaction mixture was stirred for 30 minutes while warming to room temperature. The reaction mixture was diluted with 500 mL of methylene chloride, washed with 200 mL saturated bicarbonate solution and then rewashed with water and brine. The organic phase was dried and evaporated to a flaky white
15 solid (15.7 grams). This material was used without further purification.

Mass Spectrum: m/e 286 (m+)

D. Methyl-2-methoxy-5-benzthioamido-benzoate

A 2 liter round bottom flask was charged with 15.7
20 grams (55 mmol) methyl-2-methoxy-5-benzamidobenzoate and 22.2 grams (55 mmol) of Lawesson's reagent followed by 500 mL of toluene. The reaction mixture was heated under reflux for 1.5 hours. The mixture was diluted with 500 mL of ethyl acetate, and washed with 250 mL of saturated aqueous
25 bicarbonate solution, followed by washes with water and brine. The organic layer was dried over sodium sulfate, filtered and evaporated. The residue was chromatographed on silica eluting with 96/4 methylene chloride - ether. There was obtained 14.8 grams (89%) of desired material.

30 E. Methyl-6-methoxy-2-phenyl-5-benzothiazolcarboxylate

A 2 liter round bottom flask was charged with 81 grams (246 mmol) potassium ferricyanide in 300 mL of water, the solution was warmed to 55°C. A solution of 14.78 grams (49
35 mmol) methyl-2-methoxy-5-benzthioamido-benzoate in 200 mL of methanol and 100 mL aqueous sodium hydroxide solution [13.7 gm (343 mmol) NaOH/100 mL] was then added. A yellow

-61-

precipitate was soon deposited. The mixture was stirred at 60°C for 1.5 hours. The reaction mixture was partitioned between 1 liter ethyl acetate and 300 mL 3.4 N HCl (aqueous). The organic phase was washed with water and
5 brine solution and then dried over sodium sulfate. After filtration, the filtrate was evaporated and the residue, as a mixture of carboxylates, was used directly in the next step. (affords 12.9 grams crude)

F. 6-Methoxy-2-phenyl-5-hydroxymethylbenzothiazole

10 A 1 liter round bottom flask was charged with 12.88 grams (45.2 mmol) of methyl-6-methoxy-2-phenyl-5-benzothiazolecarboxylate (as part of a mixture of regioisomeric carboxylates) in 600 mL of dry tetrahydrofuran (THF). This solution was cooled to 0°C and treated with
15 67.8 mL (67.8 mmol) of a 1.0 M solution of lithium aluminum hydride in THF. The dark reaction mixture was stirred for 5 hours under nitrogen. The reaction was worked up by the careful dropwise addition of a saturated aqueous solution of sodium sulfate until a granular precipitate formed. The
20 suspension was filtered and evaporated. The residue was dissolved in ethyl acetate and washed with water and saturated brine solution. The organic phase was dried over sodium sulfate and evaporated. This product was used directly in the next step. (10.94 grams crude yield)

25 G. 6-Methoxy-2-phenyl-benzothiazol-5-aldehyde

A solution of 10.94 grams (40.4 mmol) 6-methoxy-2-phenyl-5-hydroxymethylbenzothiazole (as part of a mixture of regioisomeric alcohols) in 600 mL of methylisobutyl ketone was treated with 42 grams (484 mmol) of manganese dioxide.
30 The mixture was heated under reflux for 1.5 hours and was then filtered through Celite®. The filtrate was evaporated in vacuo and the residue was chromatographed on silica (elution with 9/1 hexane/ethyl acetate). Mixed fractions were rechromatographed on silica using 95/5 hexane/ethyl
35 acetate to afford a total of 2.43 grams of the desired aldehyde.

Mass Spectrum: m/e 270 (m+1)

-62-

H. (2S, 3S)-(6-Methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenyl-piperidin-3-yl)-amine

A round bottom flask equipped with a Dean Stark water separator and condenser was charged with 0.96 grams (3.6 mmol) 6-methoxy-2-phenyl-benzothiazol-5-aldehyde and 0.57 grams (3.2 mmol) of (2S, 3S)-2-phenyl-3-amino-piperidine in 90 mL of hours. The reaction mixture was evaporated in vacuo and the residue was redissolved in 90 mL of 1,2-dichlorethane. The solution was treated with 0.89 grams (4.2 mmol) of sodium triacetoxyborohydride and was stirred for 16 hours at room temperature. The reaction mixture was diluted with methylene chloride (600 mL) and washed with 150 mL of bicarbonate solution, 150 mL of water and finally with 150 mL of saturated brine solution. The organic phase was dried over sodium sulfate, filtered and evaporated. The residue was chromatographed on silica gel eluting with 97/2/1 methylene chloride, methanol, conc. aqueous ammonium hydroxide solution. Product containing fractions were combined, evaporated and crystallized from hot ether - MeOH to afford 1.1 grams (80%) of the desired product.

Mass Spectrum: m/e 430 (m+1)

I. (2S, 3S)-(6-Methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenyl-piperidin-3-yl)-amine dihydrochloride salt

To a flame dried flask was added 20 mL of methanol and 1.97 mL (27.4 mmol) of acetyl chloride by slow dropwise addition. The solution was stirred for 10 min. at room temperature. A solution of (2S, 3S)-(6-methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenyl-piperidin-3-yl)-amine 2.35 grams (5.4 mmol) in methanol (prepared by warming) was then added to the reaction mixture prepared above. The resulting solution was stirred for 10 min. and then evaporated in vacuo. The residue was taken up in 500 mL of methanol, heated under reflux and filtered. The resulting solution was treated with 200 mL of diethyl ether and was allowed to stand at 5°C to effect crystallization. There was obtained 2.55 grams of the hydrochloride salt (94%).

-63-

Analysis: Calc'd for $C_{26}H_{27}N_3OS \cdot 2 HCl$: C, 62.15; H, 5.82; N, 8.36. Found: C, 62.17; H, 5.89; N, 8.31.

The title compounds of Examples 21-25 were prepared using procedures analogous to that of Example 20.

5

EXAMPLE 21

(2S,3S)-(6-Methoxy-2-cyclopropyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine

Mass Spectrum: m/e 394 (m+1)

EXAMPLE 22

10

(2S,3S)-(6-Methoxy-2-tert-butyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine

HRMS calc'd for $C_{24}H_{31}N_3OS$: 409.2181. Found: 409.22240.

EXAMPLE 23

15 (2S,3S)-(6-Isopropoxyoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine

HRMS calc'd for $C_{28}H_{31}N_3OS$: 457.2181. Found: 457.21684.

EXAMPLE 24

(2S,3S)-(6-Isopropoxyoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine

20 HRMS calc'd for $C_{23}H_{29}N_3OS$: 395.2025. Found: 395.20059.

EXAMPLE 25



(2S,3S)-(6-Trifluoromethoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine

25 HRMS calc'd for $C_{21}H_{22}F_3N_3OS$: 421.1431. Found: 421.14432.

-64-

CLAIMS

1. A compound of the formula



- 15 wherein A is a ring system selected from phenyl, naphthyl, thienyl, quinoliny and indoliny, and wherein the side chain containing NR^2R^3 is attached to a carbon atom of ring system A;

20 W is hydrogen, $(\text{C}_1\text{-C}_6)$ alkyl optionally substituted with from one to three fluorine atoms, $-\text{S}(\text{O})_v-(\text{C}_1\text{-C}_6)$ alkyl wherein v is zero, one or two, halo, benzyloxy or $(\text{C}_1\text{-C}_6)$ alkoxy optionally substituted with from one to three fluorine atoms;

25 R^1 is a 4, 5 or 6 membered heterocyclic ring containing from one to three heteroatoms selected from oxygen, nitrogen and sulfur (e.g., thiazolyl, azetidiny, pyrrolyl, pyrazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, isothiazolyl, imidazolyl, isoxazolyl, oxazolyl, pyridyl, pyrimidinyl, pyrazolyl or thiophenyl), wherein said heterocyclic ring may
30 contain from zero to three double bonds and may optionally be substituted with one or more substituents, preferably one or two substituents, independently selected from $(\text{C}_1\text{-C}_6)$ alkyl optionally substituted with from one to three fluorine atoms and $(\text{C}_1\text{-C}_6)$ alkoxy optionally substituted with from one
35 to three fluorine atoms;

the dotted lines in formula Ib indicate that one of the $\text{X}'\text{-Y}'$ and $\text{Y}'\text{-Z}'$ bonds may optionally be a double bond;

-65-

X' is selected from =CH-, -CH₂-, -O-, -S-, -SO-, -SO₂-,
-N(R⁴)-, -NH-, =N-, -CH[(C₁-C₆)alkyl]-, =C[(C₁-C₆)alkyl]-,
-CH(C₆H₅)- and =C(C₆H₅)-;

Y' is selected from C=O, C=NR⁴, C=S, =CH-, -CH₂-, =C[(C₁-
5 C₆)alkyl]-, -CH[(C₁-C₆)alkyl]-, =C(C₆H₅)-, -CH(C₆H₅)-, =N-,
-NH-, -N(R⁴)-, =C(halo)-, =C(OR⁴)-, =C(SR⁴)-, =C(NR⁴)-, -O-,
=C(CF₃)-, =C(CH₂C₆H₅)-, -S- and SO₂, wherein the phenyl
moieties of said =C(C₆H₅)- and -CH(C₆H₅)- may optionally be
substituted with from one to three substituents,
10 independently selected from halo and trifluoromethyl, and
wherein the alkyl moieties of said =[(C₁-C₆)alkyl]- and -
[CH(C₁-C₆)alkyl]- may optionally be substituted with from one
to three fluorine atoms;

Z' is selected from =CH-, -CH₂-, =N-, -NH-, -S-,
15 -N(R⁴)-, =C(C₆H₅)-, -CH(C₆H₅)-, =C[(C₁-C₆)alkyl]- and -CH[(C₁-
C₆)alkyl]-;


or X', Y' and Z', together with the two carbon atoms
shared between the benzo ring and the X'Y'Z' ring, form a
fused pyridine or pyrimidine ring;

20 R² is hydrogen or -CO₂(C₁-C₁₀)alkyl;


R³ is selected from

-66-

5




I




III

10




IV




V

15




VI

20




VII

25



VIII

or



IX

35

-67-

wherein R^6 and R^{10} are independently selected from furyl, thienyl, pyridyl, indolyl, biphenyl and phenyl, wherein said phenyl may optionally be substituted with one or two substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms, (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms, carboxy, benzyloxycarbonyl and (C_1-C_3) alkoxy-carbonyl;


R^4 is (C_1-C_6) alkyl or phenyl;

R^7 is selected from (C_3-C_4) branched alkyl, (C_5-C_6) branched alkenyl, (C_5-C_7) cycloalkyl, and the radicals named in the definition of R^6 ;

R^8 is hydrogen or (C_1-C_6) alkyl;

R^9 and R^{19} are independently selected from phenyl, biphenyl, naphthyl, pyridyl, benzhydryl, thienyl and furyl, and R^9 and R^{19} may optionally be substituted with from one to three substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms and (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms;

Y is $(CH_2)_1$, wherein 1 is an integer from one to three, or Y is a group of the formula



(J)

;

Z is oxygen, sulfur, amino, (C_1-C_3) alkylamino or $(CH_2)_n$, wherein n is zero, one or two;

x is zero, one or two;

y is zero, one or two;

z is three, four or five;

o is two or three;

p is zero or one;

r is one, two or three;

-68-

the ring containing $(CH_2)_2$ may contain from zero to three double bonds, and one of the carbon atoms of $(CH_2)_2$ may optionally be replaced by oxygen, sulfur or nitrogen;

R^{11} is thienyl, biphenyl or phenyl optionally substituted with one or two substituents independently selected from halo, (C_1-C_{10}) alkyl optionally substituted with from one to three fluorine atoms and (C_1-C_{10}) alkoxy optionally substituted with from one to three fluorine atoms;

10 X is $(CH_2)_q$ wherein q is an integer from 1 to 6, and wherein any one of the carbon-carbon single bonds in said $(CH_2)_q$ may optionally be replaced by a carbon-carbon double bond, and wherein any one of the carbon atoms of said $(CH_2)_q$ may optionally be substituted with R^{14} , and wherein any one
15 of the carbon atoms of said $(CH_2)_q$ may optionally be substituted with R^{15} ;

m is an integer from 0 to 8, and any one of the carbon-carbon single bonds of $(CH_2)_m$, wherein both carbon atoms of such bond are bonded to each other and to another
20 carbon atom in the $(CH_2)_m$ chain, may optionally be replaced by a carbon-carbon double bond or a carbon-carbon triple bond, and any one of the carbon atoms of said $(CH_2)_m$ may optionally be substituted with R^{17} ;

R^{12} is a radical selected from hydrogen, (C_1-C_6) straight
25 or branched alkyl, (C_3-C_7) cycloalkyl wherein one of the carbon atoms may optionally be replaced by nitrogen, oxygen or sulfur; aryl selected from biphenyl, phenyl, indanyl and naphthyl; heteroaryl selected from thienyl, furyl, pyridyl, thiazolyl, isothiazolyl, oxazolyl, isoxazolyl, triazolyl, tetrazolyl and quinolyl; phenyl- (C_2-C_6) alkyl, benzhydryl and
30 benzyl, wherein the point of attachment on R^{12} is a carbon atom unless R^{12} is hydrogen, and wherein each of said aryl and heteroaryl groups and the phenyl moieties of said benzyl, phenyl- (C_2-C_6) alkyl and benzhydryl may optionally be
35 substituted with one or more substituents independently selected from halo, nitro, (C_1-C_{10}) alkyl optionally

-69-

substituted with from one to three fluorine atoms,
 (C₁-C₁₀)alkoxy optionally substituted with
 from one to three fluorine atoms, amino, hydroxy-(C₁-
 C₆)alkyl, (C₁-C₆)alkoxy-(C₁-C₆)alkyl, (C₁-C₆)-alkylamino,

5 (C₁-C₆)alkyl-O-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-(C₁-C₆)alkyl,
 10 (C₁-C₆)alkyl-C(=O)-O-, (C₁-C₆)alkyl-C(=O)-(C₁-C₆)alkyl-O-,
 15 (C₁-C₆)alkyl-C(=O)-, (C₁-C₆)alkyl-C(=O)-(C₁-C₆)alkyl-,
 20 di-(C₁-C₆)alkylamino, -C(=O)NH-(C₁-C₆)alkyl,
 (C₁-C₆)-alkyl-C(=O)NH-(C₁-C₆)alkyl, -NHCH(=O) and -NHC(=O)-(C₁-C₆)alkyl;
 and wherein one of the phenyl moieties of said benzhydryl
 25 may optionally be replaced by naphthyl, thienyl, furyl or
 pyridyl;

R¹³ is hydrogen, phenyl or (C₁-C₆)alkyl;


or R¹² and R¹³, together with the carbon to which they
 are attached, form a saturated carbocyclic ring having from
 30 3 to 7 carbon atoms wherein one of said carbon atoms that is
 neither the point of attachment of the spiro ring nor
 adjacent to it may optionally be replaced by oxygen,
 nitrogen or sulfur;

R¹⁴ and R¹⁵ are each independently selected from
 35 hydrogen, hydroxy, halo, amino, oxo (=O), cyano, hydroxy-(C₁-
 C₆)alkyl, (C₁-C₆)alkoxy-(C₁-C₆)alkyl, (C₁-C₆)alkylamino,


40 di-(C₁-C₆)alkylamino, (C₁-C₆)alkoxy, -C(=O)OH,
 (C₁-C₆)alkyl-O-C(=O)-, (C₁-C₆)alkyl-O-C(=O)-(C₁-C₆)alkyl,

45

-70-



5

10 set forth in the definition of R^{12} ;

R^{16} is NHCR^{18} , $\text{NHCH}_2\text{R}^{18}$, SO_2R^{18} , CO_2H or one of the radicals
15 set forth in any of the definitions of R^{12} , R^{14} and R^{15} ;

R^{17} is oximino ($=\text{NOH}$) or one of the radicals set forth
in any of the definitions of R^{12} , R^{14} and R^{15} ; and

R^{18} is $(C_1-C_6)\text{alkyl}$, hydrogen, phenyl or phenyl $(C_1-C_6)\text{alkyl}$;

20 with the proviso that (a) when m is 0, one of R^{16} and R^{17}
is absent and the other is hydrogen, (b) when R^3 is a group
of the formula VIII, R^{14} and R^{15} cannot be attached to the
same carbon atom, (c) when R^{14} and R^{15} are attached to the
same carbon atom, then either each of R^{14} and R^{15} is
25 independently selected from hydrogen, fluoro, $(C_1-C_6)\text{alkyl}$,
hydroxy- $(C_1-C_6)\text{alkyl}$ and $(C_1-C_6)\text{alkoxy}-(C_1-C_6)\text{alkyl}$, or R^{14} and
 R^{15} , together with the carbon to which they are attached,
form a (C_3-C_6) saturated carbocyclic ring that forms a spiro
compound with the nitrogen-containing ring to which they are
30 attached; (d) R^{12} and R^{13} cannot both be hydrogen; and (e)
when R^{14} or R^{15} is attached to a carbon atom of X or (CH_2) ,
that is adjacent to the ring nitrogen, then R^{14} or R^{15} ,
respectively, must be a substituent wherein the point of
attachment is a carbon atom;

35 or a pharmaceutically acceptable salt of such compound.

2. A compound according to claim 1, wherein the
substituents at positions "2" and "3" of the nitrogen
containing ring of R^3 are in a cis configuration.

3. A compound according to claim 1, having the
40 formula Ia wherein R^3 is a group of the formula III, VII or
IX; R^2 is hydrogen; A is phenyl or indoliny1; W is $(C_1-$

-71-

C₃)alkoxy optionally substituted with from one to five fluorine atoms; and R¹ is thiazolyl, imidazolyl, thiadiazolyl, pyrrolyl, pyridyl, pyrimidinyl, pyrazolyl, thiophenyl or oxazolyl, and R¹ may optionally be substituted
5 with one or two (C₁-C₃) alkyl moieties.

4. A compound according to claim 1, having the formula Ib, wherein R³ is a group of the formula III, VII or IX; R² is hydrogen; the fused bicyclic ring system to which W and the -CH₂NR²R³ sidechain are attached is benzoxazolyl,
10 benzimidazolyl, benzisoxazolyl, benzthiophenyl or benzthiazolyl; and W is (C₁-C₆)alkoxy optionally substituted with from one to five fluorine atoms.

5. A compound according to claim 1, wherein: (a) R³ is a group of the formula III and R⁹ is benzhydryl; (b) R³ is a
15 group of the formula VII, R¹² is phenyl, each of R¹³, R¹⁴, R¹⁵ and R¹⁶ is hydrogen, m is zero and X is -(CH₂)₃-; or (c) R³ is a group of the formula IX, r is two and R¹⁹ is benzhydryl.

6. A compound according to claim 1, having the formula Ia wherein R³ is a group of the formula III wherein
20 the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, R⁹ is benzhydryl and A is phenyl.

7. A compound according to claim 1, having the formula Ia and wherein R³ is a group of the formula VII
25 wherein R¹² and the substituent at position "3" of the nitrogen containing ring are in the cis configuration, A is phenyl, R¹² is phenyl, each of R², R¹³, R¹⁴, R¹⁵ and R¹⁶ is hydrogen, m is zero, W is methoxy or isopropoxy, X is -(CH₂)₃- and R¹ is thiazolyl, imidazolyl, pyrrolyl, oxazolyl,
30 pyridyl, pyrimidinyl, pyrazolyl, thiophenyl or thiadiazolyl.

8. A compound according to claim 1, having the formula Ib wherein R³ is a group of the formula IX wherein
the substituents at positions "2" and "3" of the nitrogen containing ring are in the cis configuration, R¹⁹ is
35 benzhydryl, r is two and the fused bicyclic ring system to

-72-


which W and the $-\text{CH}_2\text{NR}^2\text{R}^3$ sidechain are attached is benzoxazolyl, benzisoxazolyl or benzthiazolyl.

9. A compound according to claim 1, having the formula Ib wherein R^3 is a group of the formula IX, R^{19} is benzhydryl, the fused bicyclic system to which W and the $-\text{CH}_2\text{NR}^2\text{R}^3$ sidechain are attached is benzoxazolyl or benzisoxazolyl, and W is methoxy.


10. A compound according to claim 1 having the formula Ib wherein R^3 is a group of the formula VII, R^{12} is phenyl, each of R^{13} , R^{14} , R^{15} and R^{16} is hydrogen, m is zero, X is $-(\text{CH}_2)_3-$, and the fused bicyclic ring system to which W and the $-\text{CH}_2\text{NR}^2\text{R}^3$ sidechain are attached is benzthiazolyl, benzoxazolyl, or benzimidazolyl.

11. A compound according to claim 1, having the formula Ia wherein R^3 is a group of the formula VII, each of R^{13} , R^{14} , R^{15} and R^{16} is hydrogen, m is zero, X is $-(\text{CH}_2)_3-$, A is phenyl, W is methoxy, and R^1 is selected from thiazolyl, imidazolyl, thiadiazolyl, pyrrolyl, pyridyl, pyrimidinyl, pyrazolyl, thiophenyl and oxazolyl.

12. A compound of the formula




XI




XII

-73-



10 wherein W, R¹, R³, X', Y' and Z' are defined in claim 1.
 13. A compound of the formula



wherein X' is -S- or -O-, and each of Y' and Z' is, independently, =N-, =CH-, =C[(C₁-C₆)alkyl]- or =C(C₆H₅)-, wherein the alkyl moiety of said =C[(C₁-C₆)alkyl] may optionally be substituted with from one to three fluorine atoms and the phenyl moiety of said =C(C₆H₅)- may optionally be substituted with from one to three substituents independently selected from halo and trifluoromethyl, with the proviso that Y' and Z' can not both be =N-, and R²² is methyl, ethyl, n-propyl, isopropyl, t-butyl, trifluoromethyl, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl or benzyl.

14. A compound according to claim 13 that is selected from the group consisting of:

6-methoxy-2-methyl-benzoxazol-5-ylaldehyde;
 6-methoxy-2-methyl-benzothiazol-5-ylaldehyde;
 6-methoxy-3-methyl-benzo[d]isoxazol-5-ylaldehyde; and
 6-methoxy-2-phenyl-benzthiazol-5-ylaldehyde.

15. A compound according to claim 1 that is selected from the group consisting of:

(2S,3S)-[5-(2,5-dimethyl-pyrrol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;

-74-

- (2S,3S)-(6-methoxy-2-methyl-benzoxazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)-amine;
- (2S,3S)-3-[2-methoxy-5-(2-thiazolyl)benzyl]amino-2-phenylpiperidine;
- 5 (2S,3S)-3-[5-imidazolyl-2-methoxybenzyl]amino-2-phenylpiperidine;
- (2S,3S)-3-[2-methoxy-5-(2-oxopyrrolidinyl)benzyl]amino-2-phenylpiperidine;
- (2S,3S)-3-[2-methoxy-5-(4-methyl-2-thiazolyl)benzyl]-
- 10 amino-2-phenylpiperidine;
- (2S,3S)-(6-methoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-3-[2-methoxy-5-(1,2,3-thiadiazol-4-yl)benzyl]-amino-2-phenylpiperidine;
- 15 (2S,3S)-3-[2-methoxy-5-(5-oxazolyl)benzyl]amino-2-phenylpiperidine;
- (2S,3S)-(6-methoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-methoxy-2-cyclopropyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- 20 (2S,3S)-(6-methoxy-2-tert-butyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-isopropoxyoxy-2-phenyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- 25 (2S,3S)-(6-isopropoxyoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (2S,3S)-(6-trifluoromethoxy-2-methyl-benzothiazol-5-ylmethyl)-(2-phenylpiperidin-3-yl)amine;
- (1SR,2SR,3SR,4RS)-(2-benzhydryl-1-aza-
- 30 bicyclo[2.2.1]hept-3-yl)-(6-methoxy-2-methyl-benzoxazol-5-ylmethyl)amine;
- (1SR,2SR,3SR,4RS)-3-[6-methoxy-3-methylbenzisoxazol-5-yl]methylanino-2-benzhydrylazanorbornane;
- (2S,3S)-(2-methoxy-5-pyridin-2-ylbenzyl)-(2-
- 35 phenylpiperidin-3-yl)amine;
- (2S,3S)-(2-methoxy-5-pyrimidin-2-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;

-75-

(2S,3S)-(2-methoxy-5-pyridin-3-ylbenzyl)-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[2-methoxy-5-(6-methylpyridin-2-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;

5 (2S,3S)-[5-(3,5-dimethylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[2-methoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;

10 (2S,3S)-[2-isopropoxy-5-(3,4,5-trimethylpyrazol-1-yl)benzyl]-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[5-(3,5-diisopropylpyrazol-1-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine;

(2S,3S)-[5-(3,5-dimethylthiophen-2-yl)-2-methoxybenzyl]-(2-phenylpiperidin-3-yl)amine; and

15 (2S,3S)-(6-methoxy-2,3-dimethyl-benzo[b]thiophen-7-ylmethyl)-(2-phenylpiperidin-3-yl)amine.

16. A pharmaceutical composition for treating or preventing a condition selected from the group consisting of inflammatory diseases, anxiety, urinary incontinence, 20 gastrointestinal disorders, depression or dysthymic disorders, psychosis, pain, allergies, chronic obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy, addiction disorders, stress related 25 somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising an amount of a compound according to claim 1 effective in preventing or treating such condition 30 and a pharmaceutically acceptable carrier.

17. A method of treating or preventing a condition selected from the group consisting of inflammatory diseases anxiety, urinary incontinence, gastrointestinal disorders, depression or dysthymic disorders, psychosis, pain, 35 allergies, chronic obstructive airways disease, hypersensitivity disorders, vasospastic diseases, fibrosing and collagen diseases, reflex sympathetic dystrophy,

-76-

addiction disorders, stress related somatic disorders, peripheral neuropathy, neuralgia, neuropathological disorders, disorders related to immune enhancement or suppression and rheumatic diseases in a mammal, comprising
5 administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 effective in preventing or treating such condition.

18. A pharmaceutical composition for antagonizing the effects of substance P in a mammal, comprising a substance
10 P antagonizing effective amount of a compound according to claim 1 and a pharmaceutically acceptable carrier.

19. A method of antagonizing the effects of substance P in a mammal, comprising administering to said mammal a substance P antagonizing effective amount of a compound
15 according to claim 1.

20. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an
20 amount of a compound according to claim 1 effective in antagonizing the effect of substance P at its receptor site and a pharmaceutically acceptable carrier.

21. A method of treating or preventing a condition in a mammal, the treatment or prevention of which is effected
25 or facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, effective in antagonizing the effect of substance
30 P at its receptor site.

22. A pharmaceutical composition for treating or preventing a condition in a mammal, the treatment or prevention of which is effected or facilitated by a decrease in substance P mediated neurotransmission, comprising an
35 amount of a compound according to claim 1, or a pharmaceutically acceptable salt thereof, effective in

-77-

treating or preventing such condition and a pharmaceutically acceptable carrier.

23. A method of treating or preventing a condition in mammal, the treatment or prevention of which is effected or
5 facilitated by a decrease in substance P mediated neurotransmission, comprising administering to a mammal in need of such treatment or prevention an amount of a compound according to claim 1 effective in treating or preventing such condition.

A. CLASSIFICATION OF SUBJECT MATTER

IPC 5 C07D401/12 A61K31/42 A61K31/445 C07D409/12 C07D413/12
 C07D417/12 C07D487/08 C07D261/20 C07D263/56 C07D275/04
 C07D277/64 C07D277/66 C07D333/54

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 5 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,91 09844 (PFIZER) 11 July 1991 see claims 1,21,23 ---	1,16,18
X	WO,A,92 20676 (PFIZER) 26 November 1992 see claims 1,34,36 ---	1,16,18
X	WO,A,90 05525 (PFIZER) 31 May 1990 see claims 1,33,38,46 ---	1,16,18
X	WO,A,90 05729 (PFIZER) 31 May 1990 see claims 1,37 ---	1,16,18
X	WO,A,92 06079 (PFIZER) 16 April 1992 see claims 1,22 ---	1,16,18
X	WO,A,92 17449 (PFIZER) 15 October 1992 see example 56 see page 11; claim 1 ---	1,16,18
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

2 December 1993

Date of mailing of the international search report

12. 01. 94

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
 NL - 2280 HV Rijswijk
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+ 31-70) 340-3016

Authorized officer

Heywood, C

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 93/09407

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,91 18878 (PFIZER) 12 December 1991 see page 1; claim 1 ---	1,16,18
P,X	WO,A,93 01170 (PFIZER) 21 January 1993 see claims 1,3,5 ---	1,16,18
P,X	WO,A,93 00330 (PFIZER) 7 January 1993 see claims 1,7,11 ---	1,16,18
A	EP,A,0 499 313 (MERCK SHARP & DOHME) 19 August 1992 see page 3; claims 1,7 ---	1-23
A	WO,A,92 01688 (PFIZER) 6 February 1992 see claims 1,24 -----	1-23

INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 93/09407

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 17,19,21,23 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/ composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/JS 93/09407

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9109844	11-07-91	EP-A- 0436334 EP-A- 0558156	10-07-91 01-09-93
WO-A-9220676	26-11-92	AU-A- 1927592	30-12-92
WO-A-9005525	31-05-90	CA-A- 2003441 EP-A- 0409931 WO-A- 9005729 US-A- 5162339	23-05-90 30-01-91 31-05-90 10-11-92
WO-A-9005729	31-05-90	WO-A- 9005525 CA-A- 2003441 EP-A- 0409931 US-A- 5162339	31-05-90 23-05-90 30-01-91 10-11-92
WO-A-9206079	16-04-92	AU-A- 8746391 CA-A- 2089736 CN-A- 1060285 EP-A- 0550635	28-04-92 29-03-92 15-04-92 14-07-93
WO-A-9217449	15-10-92	AU-A- 1244892 CN-A- 1065264	02-11-92 14-10-92
WO-A-9118878	12-12-91	AU-A- 7770391 CN-A- 1056876 EP-A- 0532515 JP-T- 5502238	31-12-91 11-12-91 24-03-93 22-04-93
WO-A-9301170	21-01-93	US-A- 5232929	03-08-93
WO-A-9300330	07-01-93	AU-A- 2188992	25-01-93
EP-A-0499313	19-08-92	JP-A- 5078354 US-A- 5242930 CA-A- 2060949	30-03-93 07-09-93 12-08-92
WO-A-9201688	06-02-92	AU-A- 7894891 CN-A- 1058405 EP-A- 0540526	18-02-92 05-02-92 12-05-93